



## **DRAFT RISK ANALYSIS REPORT**

### **APPLICATION A383**

#### **Food derived from insect and potato leafroll virus-protected potato lines RBMT21-129, RBMT21-350, and RBMT22-82**

**Note:**

This report is the “Full Assessment” as referred to in Section 15 of the *Australia New Zealand Food Authority Act (1991)*.

Public comments are now sought before completion of a Final Risk Analysis Report (referred to as the “Inquiry” in Section 16 of the Act). See under ‘Invitation for Public Submissions’ for details.

## TABLE OF CONTENTS

<b>EXECUTIVE SUMMARY .....</b>	<b>3</b>
BACKGROUND.....	3
ISSUES ADDRESSED DURING ASSESSMENT .....	3
<i>i. Safety Evaluation</i> .....	3
<i>ii. Labelling</i> .....	3
<i>iii. Public Submissions</i> .....	4
CONCLUSION.....	4
<b>INVITATION FOR PUBLIC SUBMISSIONS.....</b>	<b>5</b>
<b>INTRODUCTION.....</b>	<b>6</b>
<b>BACKGROUND TO THE APPLICATION .....</b>	<b>6</b>
<b>PUBLIC CONSULTATION.....</b>	<b>7</b>
<b>NOTIFICATION OF THE WORLD TRADE ORGANIZATION .....</b>	<b>7</b>
<b>ISSUES ADDRESSED DURING ASSESSMENT .....</b>	<b>7</b>
1. SAFETY ASSESSMENT (ATTACHMENT 2).....	7
2. LABELLING OF FOOD DERIVED FROM INSECT AND PLRV-PROTECTED POTATOES .....	9
3. ISSUES ARISING FROM PUBLIC SUBMISSIONS .....	10
4. RISK MANAGEMENT .....	12
<b>REGULATORY IMPACT ASSESSMENT .....</b>	<b>13</b>
<b>CONCLUSIONS .....</b>	<b>13</b>
<b>DRAFT VARIATION TO THE <i>FOOD STANDARDS CODE</i> .....</b>	<b>14</b>
<b>ACKNOWLEDGEMENTS .....</b>	<b>44</b>
<b>REFERENCES.....</b>	<b>45</b>
<b>DRAFT REGULATORY IMPACT ASSESSMENT .....</b>	<b>52</b>
<b>WORLD TRADE ORGANIZATION AGREEMENTS .....</b>	<b>54</b>
<b>SUMMARY OF FIRST ROUND PUBLIC SUBMISSIONS FOR APPLICATIONS A372, A375, A378, A379, A380, A381, A382, A383, A384, A385, A386, A387 &amp; A388...57</b>	<b>57</b>
<b>GENERAL ISSUES RAISED IN PUBLIC COMMENTS .....</b>	<b>66</b>

## **EXECUTIVE SUMMARY**

### **Background**

An application was received from Monsanto Australia Ltd on 30 April 1999 for the approval of food from genetically modified (GM) potato lines RBMT21-129, RBMT21-350 and RBMT22-82. The potatoes have been genetically modified to provide protection against a range of insects, including the Colorado potato beetle (CPB), as well as against potato leafroll virus (PLRV). The potatoes are known commercially as New Leaf® Plus potatoes. This report describes the scientific assessment of the application.

### **Issues addressed during assessment**

#### *i. Safety Evaluation*

The New Leaf® Plus potatoes have been evaluated according to ANZFA's safety assessment guidelines. This involves an extensive analysis of the nature of the genetic modification together with a consideration of general safety issues, toxicological issues and nutritional issues associated with the new GM food. This approach can establish whether the food produced from the New Leaf® Plus potatoes is as safe and nutritious as food produced from non-GM varieties of potatoes.

The detailed information available on the genetic modification indicates that no unintentional changes have taken place at the molecular level and that the novel genetic material is stably inserted in the potato genome and maintained over several generations.

Data on the potential toxicity and allergenicity of the proteins encoded by the transferred genes have been reviewed and indicates that the new proteins expressed in the New Leaf® Plus potatoes are non-toxic and unlikely to have allergenic potential.

Compositional analyses demonstrate no significant differences between the New Leaf® Plus potatoes and their conventional counterparts. This constitutes further evidence that no unintentional effects have occurred as a result of the genetic modification.

The impact on human health from the potential transfer of novel genetic material to cells in the human digestive tract has also been considered. The presence of novel genetic material, including two antibiotic resistance genes, in the New Leaf® Plus potatoes is not considered to pose any additional safety concerns.

In assessing all of the above data, ANZFA has concluded that the New Leaf® Plus potatoes do not raise any public health and safety concerns.

#### *ii. Labelling*

On the basis of the data considered in the safety evaluation, the New Leaf® Plus potatoes were found to be substantially equivalent to non-GM potatoes therefore no mandatory labelling is required.

It should be noted that on 28 July 2000 the Australia New Zealand Food Standards Council agreed to a revised standard which requires labelling of food where novel DNA and/or protein is present in the final food and also where the food has altered characteristics. This requirement will come into effect 12 months after the date of gazettal and may result in changes to the way in which GM foods, including those derived from New Leaf® Plus potatoes, are labelled.

*iii. Public Submissions*

Forty-five public submissions were received in relation to this application, of which only four were supportive. Those opposing the application did so primarily on the basis that they perceive GM food to be unsafe. The food safety concerns raised in submissions have been addressed by the draft safety assessment report.

**Conclusion**

ANZFA considers that food from New Leaf® Plus potatoes is as safe for human consumption as food from other commercial potato varieties and is therefore proposing an amendment to the Australian *Food Standards Code* to give approval to such food. Based on the data submitted in the present application, food derived from the New Leaf® Plus potatoes can be regarded as substantially equivalent to food derived from non-GM potatoes therefore no mandatory labelling is required, although as noted above this may change once the new labelling provisions of Standard A18 come into effect.

ANZFA now seeks public comment on the proposed amendment to Standard A18 of the *Food Standards Code* (in accordance with the procedures described in section 17 of the *Australia New Zealand Food Authority Act 1991*).

## INVITATION FOR PUBLIC SUBMISSIONS

The Authority has completed a Draft Risk Analysis Report on this application, (referred to as the 'Full Assessment' in section 15 of the Act), which includes a draft Safety Assessment report and a draft variation to the Australian *Food Standards Code*. The Authority now seeks public comment on the draft Safety Assessment Report, the draft variation to the *Food Standard Code*, and the Regulatory Impact Assessment before preparing a Final Risk Analysis Report (referred to as the 'Inquiry' in section 16 of the Act).

Written submissions containing technical or other relevant information, which will assist the Authority in preparing the Final Risk Analysis Report for this application, are invited from interested individuals and organisations. Technical information presented should be in sufficient detail to allow independent scientific assessment.

Submissions providing more general comment and opinion are also invited. The Authority's policy on the management of submissions is available from the Standards Liaison Officer upon request.

The processes of the Authority are open to public scrutiny, and any submissions received will ordinarily be placed on the public register of the Authority and made available for inspection. If you wish any information contained in a submission to remain confidential to the Authority, you should clearly identify the sensitive information and provide justification for treating it as commercial-in-confidence. The *Australia New Zealand Food Authority Act 1991* requires the Authority to treat in confidence trade secrets relating to food and any other information relating to food, the commercial value of which would be, or could reasonably be expected to be, destroyed or diminished by disclosure.

All correspondence and submissions on this matter should be addressed to the **Project Manager - Application A383** at one of the following addresses:

Australia New Zealand Food Authority

PO Box 7186  
Canberra Mail Centre ACT 2610  
AUSTRALIA  
Tel (02) 6271 2222 Fax (02) 6271 2278  
Email [info@anzfa.gov.au](mailto:info@anzfa.gov.au)

PO Box 10559  
The Terrace WELLINGTON 6036  
NEW ZEALAND  
Tel (04) 473 9942 Fax (04) 473 9855  
Email [nz.reception@anzfa.gov.au](mailto:nz.reception@anzfa.gov.au)

Submissions should be received by the Authority by **25 October 2000**.

General queries on this matter and other Authority business can be directed to the Standards Liaison Officer at the above address or by Email on [slo@anzfa.gov.au](mailto:slo@anzfa.gov.au). Submissions should not be sent by Email as the Authority cannot guarantee receipt. Requests for more general information on the Authority can be directed to the Information Officer at the above addresses.

## INTRODUCTION

The Australia New Zealand Food Authority (ANZFA) is a bi-national statutory body responsible for making recommendations on food standards which, when approved by the Australia New Zealand Food Standards Council (ANZFS), are adopted by reference and without amendment into food law. ANZFA is currently working to establish a joint *Australia New Zealand Food Standards Code* that will apply in both countries. In the interim, a system of dual standards operates for the majority of the food standards. Standard A18 – Food Produced using Gene Technology has been accepted by New Zealand, and currently applies in both countries.

Standard A18 was adopted by ANZFS as a joint Australia/New Zealand standard in July 1998 and came into force on 13 May 1999. Under this Standard, the sale of food produced using gene technology is prohibited unless the food is included in the Table to Clause 2 of the Standard. The Standard requires that a pre-market safety assessment be conducted on all foods produced using gene technology. However, the Standard provides interim arrangements for those foods currently on the market provided that an application was accepted by ANZFA on or before 30 April 1999, that the food is lawfully permitted in a country other than Australia or New Zealand, and that ANZFS has not become aware of evidence that the food poses a significant risk to public health and safety.

## BACKGROUND TO THE APPLICATION

ANZFA received an application from Monsanto Australia Ltd on 30 April 1999 to amend the Australian *Food Standards Code* to include food produced from potato lines RBMT21-129, RBMT21-350, and RBMT22-82 in the Table to Clause 2 of Standard A18 – Food Produced using Gene Technology.

The three lines of Russet Burbank potatoes have been genetically modified to provide protection against a range of insects, including the Colorado potato beetle (CPB), as well as against potato leafroll virus (PLRV). The genetically modified potatoes are known commercially as New Leaf® Plus potatoes.

The New Leaf® Plus potatoes are protected against CPB and PLRV through the transfer of two genes — the *cry3Aa* gene from the soil bacterium *Bacillus thuringiensis* subspecies *tenebrionis* (*B.t.t.*) and the *PLRVrep* gene from PLRV.

New Leaf® Plus potatoes are not grown in Australia or New Zealand and are currently not permitted to be imported into Australia and New Zealand as fresh produce. Rather, the principle food products derived from these potato varieties are likely to be imported processed food commodities such as processed potato crisps, pre-cooked French fries, potato flour and potato starch.

Direct benefits from the New Leaf® Plus potatoes are likely to accrue mainly to the primary producer who will be able to substantially reduce costs for controlling CPB and PLRV by reducing reliance on the use of agricultural chemicals. More general benefits, however, may also flow to the community as a result of reduced primary production costs.

## **PUBLIC CONSULTATION**

ANZFA completed a Notice of Application (formally referred to as the Preliminary Assessment Report) upon receipt of the application and called for public comment on 3 November 1999. A total of 45 submissions were subsequently received. Attachment 5 contains a summary of the submissions.

## **NOTIFICATION OF THE WORLD TRADE ORGANIZATION**

During the ANZFA assessment process, comments are also sought internationally from other Members of the World Trade Organization (WTO). As Members of the WTO, Australia and New Zealand are signatories to the agreements on the Application of Sanitary and Phytosanitary Measures (SPS Agreement) and on Technological Barriers to Trade (TBT Agreements) (for further details on WTO, see Attachment 4). In some circumstances, Australia and New Zealand have an obligation to notify the WTO of changes to food standards to enable other member countries of the WTO to make comment.

As there is significant international interest in the safety of these foods, the proposed changes to Standard A18 are considered to raise potential Technical Barrier to Trade or Sanitary/Phytosanitary matters and will therefore be notified to the WTO.

## **ISSUES ADDRESSED DURING ASSESSMENT**

### **1. Safety assessment (attachment 2)**

The safety assessment was performed according to the safety assessment guidelines prepared by ANZFA<sup>1</sup> and considered the following issues: (1) the nature of the genetic modification; (2) general safety issues such as novel protein expression and the potential for transfer of novel genetic material to cells in the human digestive tract; (3) toxicological issues; and (4) nutritional issues.

#### *Nature of the genetic modification*

Five genes were transferred to the New Leaf® Plus potatoes using *Agrobacterium tumefaciens*-mediated transformation – *cry3Aa*, *PLRVrep*, *nptII*, *CP4 EPSPS*, and *aad*.

The *cry3Aa* and *PLRVrep* genes are present in all three New Leaf® Plus lines. The *nptII*, *CP4 EPSPS*, and *aad* genes, however, are not present in all New Leaf® Plus lines. Lines RBMT21-129 and RBMT21-350 contain the *nptII* gene, and line RBMT22-82 contains the *aad* and *CP4 EPSPS* genes. The *aad* gene in line RBMT22-82 does not appear to be intact.

The *cry3Aa* gene is one of several isolated from *B. thuringiensis*, which encode a group of toxins known as the *Bt* toxins. These toxins are selectively active against several groups of insects such as moths and butterflies, beetles, and flies and mosquitos. The *Bt* toxin produced by the *cry3Aa* gene is known as Cry3Aa and is selectively active against a narrow range of beetles, including CPB. When a susceptible beetle ingests Cry3Aa the toxin binds to the cells lining the insect gut causing their rupture and leading to gut paralysis. The insect stops

---

<sup>1</sup> ANZFA (1999) Guidelines for the safety assessment of foods to be included in Standard A18 – food produced using gene technology.

feeding and eventually dies. Cry3Aa produces this toxic effect by binding to specific receptors in the gut of target insects. As there are no receptors for Cry3Aa on the surface of mammalian intestinal cells, humans are not susceptible to Cry3Aa.

The *PLRVrep* gene is responsible for the production of the PLRV replicase protein. The replicase protein is an enzyme whose normal function is to copy (i.e., replicate) the genome of the virus within the plant cell. It has been found that plants can be protected from infection by a virus through the expression of one of a number of the virus genes in the plant. The exact mechanism by which the viral protection occurs is unknown.

The three other genes were all used as markers to assist in the selection of transformed cells (i.e., cells to which the gene of interest has been transferred). The *nptII* gene encodes the enzyme neomycin phosphotransferase II (NPTII) and confers resistance to the antibiotics neomycin, kanamycin, and geneticin (G418). The *aad* gene encodes the enzyme streptomycin adenylyltransferase and confers resistance to the antibiotics spectinomycin and streptomycin. The *CP4 EPSPS* gene encodes the enzyme 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS) and confers resistance to the herbicide glyphosate. The glyphosate-tolerant trait of the New Leaf® Plus line RBMT22-82 was included for selection purposes only. Line RBMT22-82 is not marketed as a herbicide-tolerant potato and the trait is not utilised in the field.

Where present, all transferred genes appear to be stably integrated and maintained in the potato plants over multiple generations.

#### *General safety issues*

The potato (*Solanum tuberosum* L.) is a major food crop throughout the world and has a long history of safe use as human food. The main food products to be derived from the New Leaf® Plus potatoes will be processed food commodities such as processed potato crisps, pre-cooked French fries, potato flour and potato starch.

The New Leaf® Plus potato lines each express three of four novel proteins — Cry3Aa, the PLRV replicase, and either NPTII or CP4 EPSPS. The Cry3Aa protein is expressed in tubers at levels ranging from 0.11 - 0.9 µg protein/g fresh weight (equivalent to 0.0006 to 0.005% of the total tuber protein). The PLRV replicase protein was unable to be detected in the plants, although other evidence suggests that it is expressed. NPTII is expressed in the tuber but at levels below the limit of detection (0.3 ng protein/g fresh weight) and CP4 EPSPS is expressed in the tuber at levels ranging from 0.21 - 0.78 µg protein/g fresh weight (equivalent to 0.001 to 0.004% of the total tuber protein). The *aad* gene is not expressed in the potatoes.

The impact on human health from potential transfer of novel genetic material from New Leaf® Plus potatoes to cells in the human digestive tract was evaluated. It was concluded that transfer was extremely unlikely to occur, and unlikely to pose any special additional risks compared with the large amount of DNA naturally present in all foods. In the case of the two antibiotic resistance genes, it was concluded that even should transfer occur, the health impacts would be negligible because these antibiotic resistance genes are already commonly carried by bacteria found in the environment as well as inhabiting the human digestive tract.

### *Toxicological issues*

The levels of naturally-occurring toxins in New Leaf® Plus potatoes were assessed as well as the potential toxicity and allergenicity of the four novel proteins — Cry3Aa, PLRV replicase, CP4 EPSPS, and NPTII.

The only naturally occurring toxins in potatoes are the glycoalkaloids. The glycoalkaloid levels in the New Leaf® Plus potatoes were slightly elevated compared to the control but were within the ranges reported for commercial varieties of Russet Burbank potatoes. The slightly elevated levels of glycoalkaloids do not represent a safety concern.

The toxicity of the Cry3Aa, NPTII and CP4 EPSPS proteins has been assessed using toxicity testing in mice with no adverse findings. Toxicity testing was not considered necessary for the PLRV replicase because of the long history of exposure of human beings to this protein through the consumption of PLRV-infected potatoes. In addition, exposure to the PLRV replicase will be considerably lower from New Leaf® Plus potatoes compared to PLRV-infected potatoes. The PLRV replicase also does not share any similarity with any known toxins. On the basis of this evidence, it can be concluded that all four novel proteins are non-toxic to humans.

All four proteins are also unlikely to be allergenic to humans because none of them have the physical characteristics that are typical of allergens, share any amino acid similarity with any of the known allergens or are present in potato tubers in large quantities. Furthermore, humans have a history of exposure to all four proteins at low levels with no recorded instances of allergenicity.

### *Nutritional issues*

Detailed compositional analyses were done to establish the nutritional adequacy of the New Leaf® Plus potatoes. Constituents analysed were total solids, sugars, soluble protein, proximate (total protein, fat, crude fibre, ash, total carbohydrates and calories), amino acid, vitamin and mineral, and glycoalkaloid content. These analyses confirmed that the New Leaf® Plus potatoes are compositionally equivalent to other commercial potato varieties in terms of these constituents.

### *Conclusion*

Based on the data submitted in the present application, the New Leaf® Plus potatoes are equivalent to other commercially available potato cultivars in terms of their safety and nutritional adequacy.

## **2. Labelling of food derived from insect and PLRV-protected potatoes**

Clause 3 of Standard A18 prescribes mandatory labelling of a food produced using gene technology when it contains new or altered genetic material and where it is not substantially equivalent in any characteristic or property of the food. As the New Leaf® Plus potatoes have been found, on the basis of data submitted with the present application, to be equivalent to other commercial varieties of potatoes there is no requirement for mandatory labelling under the current standard.

It should be noted that on 28 July 2000 the Australia New Zealand Food Standards Council agreed to a revised standard which requires labelling of food where novel DNA and/or protein is present in the final food and also where the food has altered characteristics. This requirement will come into effect 12 months after the date of gazettal and may result in changes to the way in which GM foods, including those derived from New Leaf® Plus potatoes, are labelled.

### **3. Issues arising from public submissions**

#### **3.1 General issues**

Of the 45 submissions received, only a small number addressed issues specific to this application. Rather, the majority of submissions raised issues of a general nature relating to gene technology or issues that had already been addressed in the safety assessment report (see Attachment 2). A discussion of some of the general issues in relation to gene technology that were raised in public submissions can be found in Attachment 6.

#### **3.2 Specific issues**

This section of the report will only address those issues raised in public submissions that are specific to an assessment of this application.

##### *(i) Use of Bt toxins*

Mr A. Ward and the Health Department of Western Australia raised concerns about the effect of *Bt* toxin on humans. The New Zealand Ministry of Health stated that ANZFA's assessment report should explain the biochemistry of the *Bt* protein and why it is unlikely to cause any harmful effects when consumed by humans.

##### Response

The *Bt* toxins are a related group of proteins produced by different types of *B. thuringiensis* during sporulation. The *Bt* toxin being used in the New Leaf® Plus potatoes is known as Cry3Aa. Cry3Aa has been shown to be selectively active against a narrow spectrum of beetles, such as CPB.

The mode of action of the *Bt* toxin is reviewed in the draft safety assessment report and its potential toxicity to mammals is also addressed (Attachment 2). It was concluded that Cry3Aa is non-toxic to mammals, including humans. This conclusion is based on a number of different pieces of evidence.

Firstly, direct experimental evidence on the absence of toxicity in mice was provided. Doses of up to 5000 mg protein/kg body weight were used. This dose has been calculated to be at least 1.2 million times higher than estimated dietary intakes.

Secondly, the mode of action of the *Bt* toxins has been thoroughly studied. When ingested by susceptible insect species, *Bt* toxins cause lysis of midgut epithelial cells in the insect gut, which leads to gut paralysis, cessation of feeding and the eventual death of the insect. The binding of the toxin to the insect gut is essential for the onset of toxicity. Binding is mediated through specialised receptors on the cell surface. If the receptors are not present on the cell

surface, the toxin will not bind and will not be able to exert its toxic effect. This is why certain types of *Bt* toxins can be toxic to some insects and not others e.g., why Cry3Aa is toxic to some types of beetles, but not toxic to moths and butterflies. The *Bt* toxins also have not shown any ability to be able to bind to mammalian gut tissue. It can therefore be inferred from the results of these studies that the *Bt* toxins are highly unlikely to exert any toxic effects in mammals, including humans, because the cells lining the human gut lack the receptors necessary for the binding of the toxin.

Lastly, microbial preparations containing Cry3Aa have been in commercial use as an insecticide on crops such as eggplant, potatoes and tomatoes in the United States since 1989 with no reports of any adverse effects in humans.

(ii) *Estimation of dietary intakes of novel proteins*

The New Zealand Ministry of Health submitted that the insect and virus-protected potatoes are likely to be a highly consumed foodstuff and suggested that the dietary intakes of the *Bt* and viral protein present in the potatoes should be estimated.

Response

When food substances are known to be hazardous, an estimate is made of the dietary intake to determine the likely human exposure to the hazard. If exposure is likely to be low there may be less cause for concern than if exposure is likely to be high.

The *Bt* and PLRV replicase proteins are not considered hazardous, that is, they are non-toxic to mammals, including humans. Because of the absence of any hazard, an estimate of the dietary intake of both proteins was not considered essential for the safety assessment. However, it is recognised that such information may be useful in providing reassurance to the community that exposure to a novel protein is low and/or that the novel protein is likely to be present in the diet at levels well below those found to be safe in animal toxicity studies.

Cry3Aa is expressed in the New Leaf® Plus potato tubers levels ranging from 0.11 to 0.9 µg protein/g fresh weight. Therefore, if certain assumptions are made about market penetration of the New Leaf® Plus potato products, it is possible to estimate the dietary intake of the *Bt* protein.

In the case of the PLRV replicase an estimate of dietary intake is not possible as the protein cannot be detected in the New Leaf® Plus potato plants. As the PLRV replicase cannot be detected in the potatoes, it follows that dietary exposure to this protein, from the consumption of New Leaf® potatoes, will be low and certainly much lower than from virus-infected potatoes.

Australian and New Zealand consumption data is available for crisps, instant mashed potato and commercial potato fries but is not currently available for potato flour or potato starch. Excluding potato flour and starch, the average total consumption<sup>2</sup> of processed potato products per person is 19.4 g/day in Australia, and 21.5 g/day in New Zealand. If, however, the consumption figures are based only on those in the population who report consuming

---

<sup>2</sup> Calculated for all respondents

potatoes then the average total consumption is 118.2 g/day and the 95<sup>th</sup> percentile consumption is 300 g/day.

For calculation of the dietary intake of the novel proteins, the highest potato consumption figure (300 g/day) and the highest protein concentration (0.9 µg protein/g fresh weight) was used. This represents a ‘worst-case’ estimate.

To do the calculation, assumptions about the proportion of processed potato products derived from the New Leaf® Plus potatoes must be made. Data on market penetration of the New Leaf® Plus potatoes is not available. In the absence of information about market penetration, two estimates will be made — one using a very worst case estimate where it is assumed that all potato products are derived entirely from New Leaf® Plus potatoes and the other, probably more realistic estimate, where it is assumed that 10% of potato products are derived from New Leaf® Plus potatoes. The dietary intake estimates are provided in the table below:

Market penetration	Estimated dietary intake	
	µg /day	µg/kg BW/day <sup>1</sup>
100 %	270.0	4.2
10 %	27.0	0.42

<sup>1</sup> assuming a body weight of 65 kg.

The very worst-case estimate is at least 1.2 million times less than the dose found to have no adverse effects in mice (5220 mg Cry3Aa/kg BW, administered as two doses in a single day). Therefore, even if all processed potato products were to be derived from the New Leaf® Plus potatoes, a very large margin of safety exists.

The potato consumption data and estimated dietary intakes of the novel proteins are included with the draft safety assessment report (Attachment 2).

#### 4. Risk management

Under Standard A18, a GM food must undergo a safety assessment in accordance with ANZFA’s safety assessment guidelines. The requirement for the food to be labelled must also be assessed in accordance with the labelling criteria specified in Clause 3 of the standard.

On the basis the conclusions of the safety assessment report, together with a consideration of the public submissions, it is proposed that Table 1 to Clause 2 of Standard A18 be amended to include food from New Leaf® Plus potato lines RBMT21-129, RBMT21-350, and RBMT22-82. The proposed amendment is provided in Attachment 1.

In relation to labelling of the food, the safety assessment report found that, based on the data submitted in the present application, New Leaf® Plus potato lines RBMT21-129, RBMT21-350, and RBMT22-82 are substantially equivalent to other commercially available potatoes in terms of their safety and nutritional adequacy. Therefore, under the current standard, no mandatory labelling is required.

In relation to the concerns raised in the public submissions with regard to gene technology and GM food, ANZFA has prepared a public discussion paper on the safety assessment process for GM food<sup>3</sup>. This is widely available and may assist in addressing some of the concerns raised by the public. Other government and industry bodies are also addressing the broader concerns in relation to gene technology.

## **REGULATORY IMPACT ASSESSMENT**

The benefits and costs associated with the proposed amendment to Standard A18 have been analysed in a draft Regulatory Impact Assessment (Attachment 3). The benefits of the proposed Standard A18 amendment to approve food from New Leaf® Plus potatoes primarily accrue to the food industry and government, with potentially a small benefit to the consumer.

## **CONCLUSIONS**

It is concluded that:

- the introduced genes in New Leaf® Plus potato lines RBMT21-129, RBMT21-350, and RBMT22-82 are not considered to produce any additional public health and safety risk;
- based on the data submitted in the present application, New Leaf® Plus potato lines RBMT21-129, RBMT21-350, and RBMT22-82 are equivalent to other commercial varieties of potatoes in terms of their safety and nutritional adequacy;
- food derived from New Leaf® Plus potato lines RBMT21-129, RBMT21-350, and RBMT22-82 does not require labelling under the current provisions of Standard A18 as it is substantially equivalent to food derived from non-GM potatoes. Recently agreed amendments to the labelling provision of Standard A18 may result in some New Leaf® Plus potato food products being labelled in the future; and
- the benefits to government, consumers and industry associated with the proposed amendment outweigh the costs.

## **ATTACHMENTS**

1. Draft variation to the *Food Standards Code*
2. Draft safety assessment report
3. Draft regulatory impact assessment
4. World Trade Organisation Agreements
5. Summary of public comments
6. General issues raised in public comments

---

<sup>3</sup> ANZFA (2000) GM foods and the consumer: ANZFA's safety assessment process for genetically modified foods. ANZFA Occasional Paper Series No. 1.

**DRAFT VARIATION TO THE *FOOD STANDARDS CODE***

**A383 – FOOD DERIVED FROM INSECT AND POTATO LEAFROLL VIRUS-PROTECTED POTATO LINES RBMT21-129, RBMT21-350, AND RBMT22-82**

*Standard A18 is varied by inserting into Column 1 of the Table to clause 2 -*

Food derived from insect and potato leafroll virus-protected potato lines RBMT21-129, RBMT21-350, and RBMT22-82.

**DRAFT SAFETY ASSESSMENT REPORT**

**APPLICATION A383 - FOOD DERIVED FROM INSECT AND POTATO  
LEAFROLL VIRUS PROTECTED POTATO LINES RBMT21-129, RBMT21-350,  
AND RBMT22-82**

## SUMMARY AND CONCLUSIONS

### *Nature of the genetic modification*

Three lines of Russet Burbank potatoes (RBMT21-129, RBMT21-350 and RBMT22-82) were protected against Colorado potato beetle (CPB) and potato leafroll virus (PLRV) through the *Agrobacterium tumefaciens* mediated transfer of two genes — the *cry3Aa* gene from the soil bacterium *Bacillus thuringiensis* subspecies *tenebrionis* (*B.t.t.*) and the *PLRVrep* gene from PLRV. The insect and virus-protected potatoes are known as New Leaf® Plus potatoes.

The *cry3Aa* gene is responsible for the production of the Cry3Aa protein which is toxic to a narrow range of beetles, including the Colorado potato beetle. When ingested by a susceptible beetle, Cry3Aa causes lysis of midgut epithelial cells in the insect gut, leading to gut paralysis, cessation of feeding and the eventual death of the insect. Cry3Aa produces this toxic effect by binding to specific receptors in the target insects. As there are no receptors for Cry3Aa on the surface of mammalian intestinal cells, humans are not susceptible to Cry3Aa. A number of microbial pesticide products based on Cry3Aa are commercially available in the United States, with some being in use since 1989.

The *PLRVrep* gene is responsible for the production of the PLRV replicase protein. The replicase is an enzyme whose normal function is to copy (i.e., replicate) the genome of the virus within the plant cell. It has been found that plants can be protected from viral infection through the expression of one of a number of viral genes in the plant. The exact mechanism by which the viral protection occurs is unknown.

Other genes transferred to the New Leaf® Plus potatoes were the *nptII* gene (in lines RBMT21-129 and RBMT21-350), and the *CP4 EPSPS* and *aad* genes (in line RBMT22-82 only). The *nptII* and *CP4 EPSPS* genes are marker genes used for selection of transformed plant lines during the potato transformation procedure. The *nptII* gene codes for the enzyme neomycin phosphotransferase II (NPTII) and confers resistance to the antibiotics neomycin, kanamycin, and geneticin (G418). The *CP4 EPSPS* gene codes for the enzyme 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS) and confers tolerance to the herbicide, glyphosate. The glyphosate tolerant trait of the New Leaf® Plus potatoes was included for selection purposes only. New Leaf® Plus potato line RBMT22-82 is not marketed as a herbicide-tolerant potato and the trait is not utilised in the field. The *aad* gene is a marker used to select transformed bacteria from non-transformed bacteria during the DNA cloning and recombination steps undertaken in the laboratory prior to transformation of the plant cells. It codes for the enzyme streptomycin adenyltransferase, which confers resistance to the antibiotics spectinomycin and streptomycin.

The transferred genes appear to be stably integrated and both protection traits are stably maintained over multiple generations.

### *General safety issues*

The potato (*Solanum tuberosum* L.) is a major food crop throughout the world and has a long history of safe use as human food. The main food products to be derived from the New Leaf® Plus potatoes will be processed food commodities such as processed potato crisps, pre-cooked French fries, potato flour and potato starch.

The New Leaf® Plus potatoes each express three of four novel proteins — Cry3Aa, the PLRV replicase, and either NPTII or CP4 EPSPS. The Cry3Aa protein is expressed in tubers at levels ranging from 0.11 - 0.9 µg protein/g fresh weight (equivalent to 0.0006 to 0.005% of the total tuber protein). The PLRV replicase protein was unable to be detected in the plants, although other evidence suggests that it is expressed. NPTII is expressed in the tuber but at levels below the limit of detection (0.3 ng protein/g fresh weight) and CP4 EPSPS is expressed in the tuber at levels ranging from 0.21 - 0.78 µg protein/g fresh weight (equivalent to 0.001 to 0.004% of the total tuber protein).

One of the important issues to consider in relation to genetically modified foods is the impact on human health from potential transfer of novel genetic material to cells in the human digestive tract. Much of the concern in this regard is with antibiotic resistance genes. In the case of the New Leaf® Plus potatoes, it was concluded that the *nptII* and *aad* genes would be extremely unlikely to transfer to bacteria in the human digestive tract because of the number and complexity of the steps that would need to take place consecutively. More importantly, however, in the highly unlikely event that transfer did occur, the human health impacts would be negligible because both these resistance genes are already commonly found in bacteria in the environment as well as inhabiting the human digestive tract. Transfer of other novel genetic material from the New Leaf® Plus potatoes to human cells via the digestive tract was also considered to be equally unlikely. As the amount of novel genetic material in the New Leaf® Plus potatoes is minute compared to the total amount of DNA present it is unlikely to pose any special additional risks compared with the large amount of DNA naturally present in all foods.

### ***Toxicological issues***

The levels of naturally-occurring toxins in New Leaf® Plus potatoes were assessed as well as the potential toxicity and allergenicity of the four novel proteins — Cry3Aa, PLRV replicase, CP4 EPSPS, and NPTII.

The only naturally-occurring toxins in potatoes are the glycoalkaloids. The glycoalkaloid levels in the New Leaf® Plus potatoes were slightly elevated compared to the control but were at the lower end of the range reported for commercial varieties of Russet Burbank potatoes. The slightly elevated levels of glycoalkaloids are not statistically significant and do not represent a safety concern.

Acute oral toxicity testing in mice had been done previously for the Cry3Aa, NPTII and CP4 EPSPS proteins under other applications and it was concluded that all three proteins are non-toxic to humans. No additional evidence has come to light that would alter this conclusion. Dietary intake assessments indicate that exposure to all three proteins from the consumption of New Leaf® Plus potatoes will be low.

The potential toxicity of the PLRV replicase had not previously been considered. Human beings have a long history of exposure to the PLRV replicase through the consumption of PLRV-infected plants. In addition, the expression levels of the PLRV replicase are likely to be lower in New Leaf® Plus potatoes than in PLRV-infected potatoes. Therefore, populations consuming New Leaf® Plus potatoes will most likely have lower exposure levels to the PLRV replicase than they would through the consumption of PLRV-infected potatoes. There is also no significant similarity between the PLRV replicase and any known toxins.

On the basis of this evidence, it was concluded that the PLRV replicase protein, as expressed in the New Leaf® Plus potatoes, is non-toxic to humans.

In terms of the potential allergenicity of the four new proteins, it has previously been concluded that Cry3Aa, NPTII and CP4 EPSPS are unlikely to be allergenic to humans. No additional data or evidence has emerged which would necessitate revising this conclusion. The PLRV replicase protein in the New Leaf® Plus potatoes has no significant similarity to any known allergens, nor is it present in large quantities in potato tubers. On the basis of this information and the fact that humans have a long history of exposure to low levels of this protein, through the consumption of PLRV-infected potatoes, with no recorded instances of allergenicity, it can be concluded that the PLRV replicase protein is unlikely to be allergenic to humans.

### ***Nutritional issues***

Detailed compositional analyses were done to establish the nutritional adequacy of the New Leaf® Plus potatoes, and to compare them to non-modified control lines. Analyses were done of total solids, dextrose, sucrose, soluble protein, proximate (total protein, fat, crude fibre, ash, total carbohydrates and calories), amino acid, and vitamin and mineral content. Some minor differences were observed for some constituents however these were not biologically significant and the values reported were all within the literature reported ranges for the Russet Burbank cultivar. On the basis of the data submitted in the present application, the New Leaf® Plus potatoes are compositionally equivalent to other commercial Russet Burbank potato cultivars.

### ***Conclusion***

Based on the data submitted in the present application, New Leaf® Plus potato lines RBMT21-129, RBMT21-350 and RBMT22-82 are equivalent to other commercially available Russet Burbank potato cultivars in terms of their safety and nutritional adequacy.

## 1. BACKGROUND

Monsanto Australia Ltd have made an application to ANZFA to amend Standard A18 of the Australian *Food Standards Code* to include food derived from potatoes which have been genetically modified to be protected against Colorado potato beetle (*Leptinotarsa decemlineata* Say.), one of the major pests of potatoes in North America, and potato leafroll virus (PLRV), a major viral disease of potatoes worldwide. The potatoes are known commercially as New Leaf® Plus potatoes.

Protection against Colorado potato beetle is achieved through expression in the plant of the insecticidal protein, Cry3Aa. Cry3Aa is produced naturally by the *tenebrionis* subspecies of the spore-forming soil bacterium *Bacillus thuringiensis* (*B.t.t.*). The majority of described *B. thuringiensis* strains produce insecticidal proteins active against lepidopteran insects (larvae of moths and butterflies) and a few are reported to have activity against dipteran insects (mosquitos and flies). The Cry3Aa protein, however, is toxic to a narrow spectrum of coleopteran insects (beetles) and shows no activity against other groups of insects such as the lepidopterans or dipterans (Herrnstadt *et al* 1986).

Two commercially available microbial pesticide products based on *B.t.t.* (M-One® and Foil®) have been in use in the United States since 1989. In addition, a bio-insecticide known commercially as MYX 1806 comprising Cry3Aa genetically engineered into the bacterium *Pseudomonas fluorescens*, which has been rendered non-viable, has been commercially available in the United States since 1991.

PLRV is a spherical RNA virus belonging to the luteovirus group of plant viruses and is transmitted primarily by the green peach aphid (*Myzus persicae*). Protection against PLRV is achieved through the expression, in the plant, of a gene derived from the viral genome, which codes for the viral replicase protein. The replicase is an enzyme that copies (i.e., replicates) the RNA genome of the virus within the plant cell. The expression of plant viral genes in plants has been shown to confer varying degrees of protection against subsequent infection by the plant virus from which the gene was derived (reviewed in Lomonosoff 1995). The exact mechanism by which the protection is conferred is unknown.

New Leaf® Plus potatoes are not grown in Australia or New Zealand and are currently not permitted to be imported into Australia and New Zealand as fresh produce. Rather, they currently enter into the market in imported processed food commodities such as processed potato crisps, pre-cooked French fries, potato flour and potato starch.

## 2. DESCRIPTION OF THE GENETIC MODIFICATION

### 2.1 *Methods used in the genetic modification*

Russet Burbank potatoes were transformed with one of two plasmids, PV-STMT21 and PV-STMT22, using *Agrobacterium*-mediated transformation of stem sections.

### 2.2 *Function and regulation of the novel genes*

*Agrobacterium*-mediated transformation of potatoes with PV-STMT21 and PV-STMT22 resulted in the transfer of the three of four genes expression cassettes — *cry3Aa*, *PLRVrep*,

and either *nptII* or *CP4 EPSPS*. Each of these expression cassettes is described in Table 1 below.

**Table 1: Description of the gene expression cassettes in PV-STMT21 and PV-STMT22**

Cassette	Genetic element	Source	Function
<i>cry3Aa</i>	ArabSSU1A promoter	<i>Arabidopsis thaliana</i> ribulose-1,5-bisphosphate carboxylase (Rubisco) small subunit at 1A promoter (Almeida <i>et al</i> 1989, Wong <i>et al</i> 1992).	Constitutive plant promoter.
	<i>cry3Aa</i>	Coding region of the <i>B.t.t.</i> Band 3 protein (Perlak <i>et al</i> 1993).	Confers protection against a narrow spectrum of Coleopterans, including Colorado potato beetle.
	NOS 3' terminator	The 3' terminator region of the nopaline synthase gene from the Ti plasmid of <i>Agrobacterium tumefaciens</i> (Depicker <i>et al</i> 1982, Bevan <i>et al</i> 1983).	Contains signals for termination of transcription and directs polyadenylation.
<i>PLRVrep</i>	35S promoter	A promoter derived from Figwort mosaic virus (FMV) (Richins <i>et al</i> 1987) containing the soybean heatshock protein 17.9 kDa 5' 77-nucleotide leader sequence (Raschke <i>et al</i> 1988).	A promoter of high-level constitutive gene expression in plant tissues.
	<i>PLRVrep</i>	Coding region for the putative PLRV replicase protein (Van der Wilk <i>et al</i> 1989).	Replicates the RNA genome of the virus.
	E9 3'	The 3' non-translated region of the pea ribulose-1,5-bisphosphate carboxylase small subunit ( <i>rbcS</i> ) E9 gene (Coruzzi <i>et al</i> 1984).	Contains signals for termination of transcription and directs polyadenylation.
<i>nptII</i> (PV-STMT21)	P-NOS	The promoter region of the nopaline synthase gene from the Ti plasmid of <i>Agrobacterium tumefaciens</i> (Fraley <i>et al</i> 1983).	A promoter of low-level constitutive gene expression in plant tissues.
	<i>nptII</i>	The gene coding for neomycin phosphotransferase II from Tn5 in <i>Escherichia coli</i> (Beck <i>et al</i> 1982).	Confers resistance to the antibiotics kanamycin and neomycin. Used as a selectable marker for plant transformation (Horsch <i>et al</i> 1984, DeBlock <i>et al</i> 1984).
	NOS 3'	The 3' terminator region of the nopaline synthase gene from the Ti plasmid of <i>Agrobacterium tumefaciens</i> (Depicker <i>et al</i> 1982, Bevan <i>et al</i> 1983).	Contains signals for termination of transcription and directs polyadenylation.
<i>CP4 EPSPS</i> (PV-STMT22)	35S promoter	A promoter derived from Figwort mosaic virus (FMV) (Richins <i>et al</i> 1987) containing the soybean heatshock protein 17.9 kDa 5' 77-nucleotide leader sequence (Raschke <i>et al</i> 1988).	A promoter of high level constitutive gene expression in plant tissues.

CTP2	The chloroplast transit peptide from the EPSPS gene of <i>Arabidopsis thaliana</i> (Della-Cioppa <i>et al</i> 1986)	Targets proteins to the chloroplasts of higher plants.
CP4 EPSPS	The gene from <i>Agrobacterium</i> sp. strain CP4 (Barry <i>et al</i> 1992) coding for the enzyme 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS).	Confers tolerance to the herbicide, glyphosate. Used as a dominant selectable marker in tissue culture following transformation (Howe <i>et al</i> 1992).
E9 3'	The 3' non-translated region of the pea ribulose-1,5-bisphosphate carboxylase small subunit (rbcS) E9 gene (Coruzzi <i>et al</i> 1984).	Contains signals for termination of transcription and directs polyadenylation.

### The *cry3Aa* gene

The *cry3Aa* gene was isolated from the DNA of *B.t.t* strain BI 256-82 (Krieg *et al* 1983). A full length clone and complete nucleotide sequence of the *cry3Aa* gene has been published (McPherson *et al* 1988, Perlak *et al* 1993). The gene is one of several that have been isolated from *B. thuringiensis* and which encode a group of toxins known as the  $\delta$ -endotoxins or the crystal proteins. These toxins are selectively active against several Orders of insects such as the Lepidoptera, Coleoptera, and Diptera. The crystal proteins are produced by the bacterium during sporulation. The protein product of the *cry3Aa* gene, Cry3Aa, is selectively active against a narrow spectrum of Coleoptera (MacIntosh *et al* 1990). When ingested by susceptible insect species, the crystal proteins cause lysis of midgut epithelial cells in the insect gut, which leads to gut paralysis, cessation of feeding and the eventual death of the insect (Höfte and Whiteley 1989). Cytolytic effects on the midgut cells are mediated by binding of the activated toxin to specialised receptors on the cell surface. This binding of the toxin to specialised receptors has been shown to be essential for the onset of toxicity (Wolfersberger 1990, Ferré *et al* 1991). Following binding of activated toxin to the receptors, a rapid change in permeability of midgut cells is observed where there is an influx of ions and water in the cell, resulting in its eventual lysis (Knowles and Ellar 1987).

### The *PLRVrep* gene

The *PLRVrep* gene was isolated from PLRV isolate LR-7 (Kaniewski *et al* 1995). The genome of PLRV has been fully sequenced (Mayo *et al* 1989). The *PLRVrep* gene consists of two overlapping open reading frames, which together are thought to encode the PLRV replicase (Miller *et al* 1995, Murphy *et al* 1995, Van der Wilk *et al* 1989).

The strategy of expressing viral genes in plants to confer protection against an infecting virus is known as pathogen-derived resistance. Sanford and Johnson (1985) first developed pathogen-derived resistance as a theoretical concept, when they proposed that resistance genes against a pathogen could be derived from the genome of the pathogen itself. This approach was first successfully applied against tobacco mosaic virus (TMV) where disease development was found to be delayed in TMV inoculated plants expressing the TMV coat protein gene (Powell *et al* 1986). The exact mechanism by which the protection occurs is unknown.

### The *nptII* gene

The *nptII* gene is widely used as a selectable marker in the transformation of plants (Kärenlampi 1996). The gene functions as a dominant selectable marker in the initial, laboratory stages of plant cell selection following transformation (Horsch *et al* 1984, DeBlock *et al* 1984). It codes for the enzyme neomycin phosphotransferase II (NPTII) and confers resistance to the aminoglycoside antibiotics, neomycin, kanamycin, and geneticin (G418). The *nptII* gene is transferred along with the *cry3Aa* and *PLRVrep* genes, enabling those plant cells successfully transformed with the *cry3Aa* and *PLRVrep* genes to grow in the presence of kanamycin. Those cells that lack the *nptII* gene, and hence the *cry3Aa* and *PLRVrep* genes, will not grow and divide in the presence of kanamycin.

### The *CP4 EPSPS* gene

The *CP4 EPSPS* gene was isolated from *Agrobacterium* sp. strain CP4 (Barry *et al* 1992). The *CP4 EPSPS* gene codes for the enzyme 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS) and confers resistance to the herbicide, glyphosate. Plant EPSPS is inhibited by glyphosate and plant cells treated with glyphosate are unable to synthesize the aromatic amino acids essential for protein synthesis resulting in the death of the plant cells (Barry *et al* 1992). In contrast, the *CP4 EPSPS* is not inhibited by glyphosate. Thus when the *CP4 EPSPS* gene is transferred along with the *cry3Aa* and *PLRVrep* genes it enables those plant cells successfully transformed with the *cry3Aa* and *PLRVrep* genes to grow in the presence of glyphosate. Those cells that lack the *CP4 EPSPS* gene, and hence the *cry3Aa* and *PLRVrep* genes, will not grow and divide in the presence of glyphosate.

### Other genetic elements

The plasmid vectors, PV-STMT21 and PV-STMT22, are double border binary plant transformation vectors, which differ only in the plant cell selectable marker region. Both plasmid vectors contain well characterised DNA segments required for selection and replication of the plasmids in bacteria as well as the right and left borders delineating the region of DNA (T-DNA) which is transferred into the plant genomic DNA. This is the region into which the gene of interest, and the plant cell selectable marker, is inserted. DNA residing outside the T-DNA region does not normally get transferred into plant genomic DNA (Zambryski 1992).

The genetic elements are described in Table 2 below.

**Table 2: Description of other genetic elements contained within PV-STMT21 and PV-STMT22**

Genetic element	Source	Function
<b><i>aad</i> (resides outside the T-DNA)</b>	Gene coding for streptomycin adenylyltransferase from transposon Tn7 in <i>Escherichia coli</i> (Fling <i>et al</i> 1985).	Confers resistance to the antibiotics spectinomycin and streptomycin.
<b>LB</b>	A 0.45 kb fragment of the octopine Ti plasmid pTi5955, which contains the 24 bp T-DNA left border (LB) region (Barker <i>et al</i> 1983).	Terminates the transfer of the T-DNA from <i>A. tumefaciens</i> to the plant genome.
<b><i>oriV</i> (resides outside the T-DNA region)</b>	A 1.3 kb origin of replication region derived from the broad-host range RK2 plasmid of <i>Agrobacterium</i> (Stalker <i>et al</i> 1981).	Allows plasmids to replicate in <i>A. tumefaciens</i> .

<b>ori-322/rop region (resides outside the T-DNA region)</b>	A 1.8 kb segment of the plasmid pBR322 which contains the origin of replication region and the <i>bom</i> site for the conjugational transfer.	Allows for maintenance of plasmids in <i>E. coli</i> and their conjugal transfer into <i>A. tumefaciens</i> cells (Bolivar <i>et al</i> 1977, Sutcliffe 1978).
<b>RB</b>	A 0.36 kb fragment from the pTiT37 plasmid containing the 24 bp nopaline-type T-DNA right border (RB) region. (Depicker <i>et al</i> 1982).	The RB region is used to initiate T-DNA transfer from <i>A. tumefaciens</i> to the plant genome.

---

The *aad* gene is used as a marker to select transformed bacteria from non-transformed bacteria during the DNA cloning and recombination steps undertaken in the laboratory prior to transformation of the plant cells. Only those bacterial cells that have been transformed with the plasmid containing the *aad* gene, and hence the genes of interest (in this case the *cry3Aa* and *PLRVrep* genes) will grow. The *aad* gene is under the control of a bacterial promoter and would therefore not be expressed in transformed plant cells.

The host for all DNA cloning and vector construction was *E. coli* strain MV1190, a derivative of the common laboratory *E. coli* K-12 strain (Bachmann 1987).

### 2.3 Characterisation of the genes in the plant

Studies submitted by Monsanto:

Rochester, D.E. and Lavrik, P.B. (1996). Molecular characterisation of potato leafroll virus and Colorado potato beetle resistant Russet Burbank potato lines transformed with plant vector PV-STMT21. Monsanto Study No. 96-01-37-20.

Rochester, D.E. and Lavrik, P.B. (1996). Molecular characterisation of potato leafroll virus and Colorado potato beetle resistant Russet Burbank potato lines transformed with plant vector PV-STMT22. Monsanto Study No. 96-01-37-22.

Reiser, S.E. *et al* (1999). Characterisation of T-DNA inserts present in New Leaf® Plus potato line number RBMT21-350 by Southern blot analysis. Monsanto Study No. 98-01-37-31.

Krohn, B.M. *et al* (1999). Characterisation of T-DNA inserts present in New Leaf® Plus potato line number RBMT22-082 by Southern blot analysis. Monsanto Study No. 98-01-37-32.

Rogan, G.J. *et al* (1999). Characterisation of T-DNA inserts present in New Leaf® Plus potato line number RBMT21-129 by Southern blot analysis. Monsanto Study No. 98-01-37-30.

Seven lines of transformed Russet Burbank potatoes were produced but only three lines have been commercialised as New Leaf® Plus potatoes. Two of the potato lines, RBMT21-129 and RBMT21-350 were transformed with plasmid PV-STMT21, which contains the *nptII* gene as a selectable marker and the third potato line, RBMT22-82, was transformed with plasmid PV-STMT22 which contains the *CP4 EPSPS* gene as the selectable marker.

The transferred genes in the New Leaf® potatoes were characterised using the polymerase chain reaction (PCR) technique (Saiki *et al* 1985). PCR is a method whereby a particular segment of DNA between two known sequences can be specifically amplified using a pair of complementary single stranded DNA primers. The PCR product can be visualised and analysed using agarose gel electrophoresis by comparison against appropriate molecular size markers and positive controls.

Genomic DNA was isolated from young leaf tissue from potato lines RBMT21-129, RBMT21-350 and RBMT22-82 and a control Russet Burbank line which were collected from a 1995 field trial in the United States. PCR was used to obtain the following information: (1) the identity of the genetic elements inserted into the genomic DNA of each of the potato lines; (2) the integrity of the inserted T-DNA; and (3) whether there had been any transfer of sequences residing outside of the T-DNA region of the plant transformation vectors (e.g. the *aad* gene and the *oriV* region).


PCR primer pairs were designed to detect the presence of each of the transferred genes by specifically amplifying their coding regions. Different sets of primers were also designed to ascertain whether the integrity of the linkage between the three transferred genetic elements had been maintained during the T-DNA transfer. Primers were also designed to detect the presence of the *aad* gene and the *oriV* region, both of which reside outside the T-DNA region. PCR products obtained from these reactions using genomic DNA from the transformed potatoes were compared to PCR products obtained from purified plasmid DNA.

The results of the PCR analysis confirm that potato lines RBMT21-129 and RBMT21-350 contain intact *PLRVrep*, *cry3Aa* and *nptII* genes and that line RBMT22-82 contains intact *PLRVrep*, *cry3Aa* and *CP4 EPSPS* genes. The PCR analysis did not detect the transfer of either the *aad* gene or the *oriV* region in any of the lines.


The applicant also used Southern blotting to characterise the T-DNA insertion into the potato lines and to confirm the results of the PCR. Southern blotting is a sensitive technique that enables the detection of specific sequences among DNA fragments separated using gel electrophoresis (Southern 1975). The overall pattern of the specific fragments detected can be used to characterise the nature of the T-DNA insertion into the genome (e.g. how many sites in the genome has the T-DNA have inserted into, whether the inserted copies are intact, etc).

The results of the Southern blotting indicate the following for each of the New Leaf® potato lines:


- (i) line RBMT21-129 — insertion of the T-DNA occurred at two sites. One of the insertion events begins at the right border of the T-DNA and continues through the *PLRVrep* gene cassette, the *cry3Aa* gene cassette and the *nptII* coding region, terminating within the NOS promoter. This T-DNA insertion was not completely resolved at the left border resulting in the partial deletion of the 5' end of the NOS promoter used to express the *nptII* gene. The second insert consists of the *PLRVrep* gene and a partially deleted *cry3Aa* gene cassette. The ArabSSU1A promoter of the *cry3Aa* gene, as well as a portion of the 5' coding region of the *cry3Aa* gene, were deleted upon integration into the plant genome. The partial *cry3Aa* gene is still associated with its NOS 3' terminator region. Northern and western blot analysis was unable to detect any protein or mRNA from the truncated *cry3Aa* gene. This T-DNA insertion was not completely resolved at the right border, resulting in the deletion of the 35S promoter as well as a portion of the 5' end of the *PLRVrep* gene. Northern blot analysis confirmed that the partial *PLRVrep* gene does not give rise to any detectable mRNA. A map of the two insertion sites is given below:



- (ii) line RBMT21-350 — insertion of the T-DNA occurred at two sites. At one site, intact copies of all three genes had been inserted. At the second site, a less than full-length copy of the T-DNA had been inserted resulting in a truncated copy of the *PLRVrep* gene, lacking the 35S promoter region. Northern blot analysis demonstrated that no detectable RNA transcript was produced from the truncated *PLRVrep* gene. The remaining two genes were intact. A map of the two insertion sites is given below:



- (iii) line RBMT22-82 — insertion of the T-DNA occurred at three sites. All three copies of the T-DNA contain intact coding regions for the *PLRVrep* gene and the *cry3Aa* gene. Two copies of the T-DNA contain an intact coding region of the *CP4 EPSPS* gene. At one site, however, a less than full-length copy of the *CP4 EPSPS* gene had been inserted. Northern analysis was not able to detect any mRNA being produced from the truncated *CP4 EPSPS* gene suggesting that, as the promoter region for the truncated *CP4 EPSPS* gene is intact, mRNA does not accumulate to any detectable levels. For another T-DNA, DNA sequence beyond the RB was also inserted. This DNA is adjoined to the RB of the T-DNA and contains the *aad* gene and the *ori-322* region. This result conflicts with that of the PCR analyses, which were unable to detect the *aad* gene. The failure to detect the *aad* gene by PCR suggests that the gene is probably not intact. A map of the three insertion sites is given below:



## Conclusion

The genetic elements transferred to each of the New Leaf® Plus lines are summarised in Table 4 below.

**Table 3: Intact genetic elements present in the New Leaf® Plus potatoes**

Plant line	<i>PLRVrep</i>	<i>cry3Aa</i>	<i>nptII</i>	<i>CP4 EPSPS</i>	<i>ori-322</i>	<i>aad</i>
RBMT21-129	√	√	√	-	-	-
RBMT21-350	√	√	√	-	-	-
RBMT22-82	√	√	-	√	√	√ (probably not intact)

### 2.4 Stability of the genetic changes

The New Leaf® Plus potatoes have been planted in field trials since 1994. The applicant reports that the potato lines, which are the subject of this application, were selected on the basis that both the PLRV and Colorado potato beetle protection traits were stable throughout multiple generations of vegetative propagation of potato for seed production.

## Conclusion

The *cry3Aa* gene and the *PLRVrep* gene in the New Leaf® Plus potato lines appear to be stably integrated with both traits being stably maintained over multiple generations.

## 3. GENERAL SAFETY ISSUES

The New Leaf® Plus potatoes have been assessed according to ANZFA's safety assessment guidelines relating to Group D foods, i.e., plants or animals that contain new or altered genetic material (ANZFA 1999).

### 3.1 *History of use*

The potato (*Solanum tuberosum* L.) is a major food crop throughout the world (Simmonds 1976). It was introduced into Europe from South America in the 16<sup>th</sup> century and is cultivated for the production of underground tubers.

Potatoes are generally consumed either cooked (as a fresh vegetable) or processed into crisps, potato flour or potato starch. They are rarely consumed raw because of the indigestibility of ungelatinised potato starch and the presence of protease inhibitors (Burton 1989).

### 3.2 *Nature of novel protein*

#### **Cry3Aa**

Cry3Aa is a protein of 644 amino acids (molecular mass 73 kDa), which is produced by *B. thuringiensis* during sporulation and is encoded by the *cry3Aa* gene. The *cry3Aa* gene was isolated from *B. thuringiensis* subsp. *tenebrionis* (*B.t.t*) strain BI 256-82. In addition to the full length Cry3Aa protein, *B.t.t* also produces a smaller form of the protein known as *B.t.t* band 3 (McPherson *et al* 1988). *B.t.t* band 3 has a molecular weight of 68 kDa (597 amino acids) and results from an internal translation initiation event within the same gene starting at an internal methionine codon at amino acid position 48. This protein has been shown to possess the same insecticidal activity and selectivity to Colorado potato beetle larvae as the full-length Cry3Aa.

The gene encoding *B.t.t* band 3 protein was engineered for plant expression by being completely re-synthesised to substitute the existing bacteria-preferred codons with plant-preferred codons (Perlak *et al* 1993). The genetic code is degenerate meaning that a given amino acid may be specified by more than one codon. For example, four different codons can be used to specify alanine. It has been found that plants often prefer different codons to bacteria to specify the same amino acid, and this can affect the expression levels of bacterial genes when they are transferred to plant cells. It has been shown that the plant expression of bacterial genes can be improved if the bacteria-preferred codons are substituted with plant-preferred codons (Perlak *et al* 1990). The re-synthesis of the gene encoding the band 3 protein, to substitute plant-preferred codons for bacteria-preferred codons, changed 399 out of 1791 nucleotides without altering the amino acid sequence. The re-synthesised *cry3Aa* gene therefore expresses a protein that is identical to that produced by *B. thuringiensis* subsp. *tenebrionis*.

#### **PLRV replicase protein**

The PLRV replicase protein has a molecular mass of 130 kDa and is encoded by two overlapping open reading frames, ORF 1 and ORF 2 (Miller *et al* 1995, Murphy *et al* 1995, Van der Wilk *et al* 1989). ORF 1 encodes a genome-linked protein (VPg) (Van der Wilk *et al* 1997) and motifs characteristic of serine-like proteases (Gorbalenya *et al* 1989, Bazan and Fletterick 1989). The VPg is covalently bound to the 5' end of the PLRV genome and is thought to be important in the initiation of viral replication. The VPg and the serine protease are produced in equimolar amounts. The serine protease is an endoprotease that is thought to be responsible for cleaving the ORF1/ORF2 gene product into its functional components. This would be required to release the VPg and make it available to bind to the viral genome. The serine protease is not packaged into the PLRV virion. ORF 2, which overlaps ORF1 is

expressed through a –1 translational frameshift at a ratio of approximately 99:1 to ORF1 (Prüfer *et al* 1992). ORF2 encodes the putative RNA dependent RNA polymerase (Van der Wilk *et al* 1989, Mayo *et al* 1989, Habili and Symons 1989). The role of the replicase is to replicate the PLRV RNA genome. The replicase is not packaged into the PLRV virion.

### **Neomycin phosphotransferase II**

Neomycin phosphotransferase II (NPTII) is an enzyme with a molecular mass of 29 kDa that catalyses the transfer of a phosphate group from adenosine 5'-triphosphate (ATP) to a hydroxyl group of aminoglycoside antibiotics, including neomycin, kanamycin and gentamicin A and B, thereby inactivating the antibiotics (Davies *et al* 1986). The enzyme is encoded by the *nptII* gene, which is derived from transposon Tn5 from the bacterium *E. coli* (Beck *et al* 1982).

### **CP4 5-enolpyruvylshikimate-3-phosphate synthase**

CP4 5-enolpyruvylshikimate-3-phosphate synthase (CP4 EPSPS) is an enzyme with a molecular weight of 48 kDa and is part of the shikimate metabolic pathway for the biosynthesis of aromatic amino acids in plants, bacteria and fungi (Levin and Springson 1964). The aromatic amino acid pathway is not present in mammalian cells (Cole 1985). The CP4 EPSPS enzyme is targeted to the chloroplast using a chloroplast transit peptide sequence derived from the *Arabidopsis thaliana* EPSPS (CTP 2). The aromatic amino acid biosynthetic pathway and endogenous EPSPS activity is located in the chloroplast. *In vitro* chloroplast uptake assays have shown that the *A. thaliana* EPSPS CTP delivers CP4 EPSPS to the chloroplast and is subsequently cleaved from the pre-protein, yielding mature CP4 EPSPS with no CTP amino acids retained (della Ciopa *et al* 1986).

### **3.3 Expression of novel protein in the plant**

Studies submitted by Monsanto:

Lavrik, P.B. (1997). Registration of the plant pesticide potato leafroll virus (PLRV) *orf1/orf2* gene: product chemistry. Monsanto Study No. 97-094E.

Lavrik, P.B. and Grace, A.M. (1996). Expression levels of *B.t.t.*, NPTII and CP4 EPSPS proteins in tissues derived from Russet Burbank potato plants resistant to Colorado potato beetle and potato leafroll virus. Monsanto Study No. 96-01-37-21.

Reding, H.K. (1998). Safety of PLRV replicase in NewLeaf® Plus potatoes. Monsanto Company. EPA application No. 198176.

The PLRV replicase could not be detected using Western blot analysis in any of the transformed plant lines. This is despite the fact that an mRNA of the correct size (3.8 kb) could be detected, demonstrating that the transferred *PLRVrep* gene is transcribed *in vivo*. Therefore, there is no direct evidence that any replicase protein is produced in the New Leaf® potatoes. The applicant, however, reports indirect evidence for the expression of the replicase protein from a separate set of studies where a frameshift mutation was introduced into the *PLRVrep* gene and subsequently transferred to potatoes (Kaniewski *et al* 1995). The frameshift mutation would result in the translation of a non-functional protein. Transgenic plants containing the mutated *PLRVrep* gene are no longer resistant to infection by PLRV suggesting that protection against PLRV is protein-mediated. These data taken together suggest that the PLRV replicase is most likely expressed in the New Leaf® Plus potatoes but

at a level or in a form which is unable to be detected using standard methods. It should also be noted that the PLRV replicase could not be detected in PLRV-infected plants (Van der Wilk *et al* 1997). It is speculated that the failure to detect the PLRV replicase in both virus infected and New Leaf® Plus potatoes expressing the *PLRVrep* gene is due to self-proteolysis mediated by a putative serine endoprotease domain contained within the protein.

The expression levels of the other three proteins, Cry3Aa, NPTII and CP4 EPSPS, in leaf and tuber tissues of transformed potato lines RBMT21-129, RBMT21-350, and RBMT22-82 were determined using enzyme linked immunosorbent assay (ELISA). ELISA is a technique that uses highly specific antibodies to identify proteins. The assay system is capable of quantifying proteins in crude tissue extracts.

Tissues were obtained from potato plants grown in field trials during 1995 at three locations in the Northwest region of the USA. A non-transformed Russet Burbank line was used as the control. At all locations, the field trials were arranged in eight replicate randomised complete block design. Leaf samples were collected approximately 10 weeks post planting. The first fully expanded compound leaf from each of four or six plants within each plot (replicate) were collected. Tuber samples were collected at harvest from each of the field sites. Ten tubers per plot from four of the eight plots were collected. All extracts were analysed in triplicate. The results are summarised in Table 4 below.

**Table 4: Protein expression levels in tissue of Russet Burbank lines transformed with either PV-STM21 or PV-STM22 – multi site data.**

Plant line	Tissue	Protein Expression level ( $\mu\text{g protein/g fresh weight}$ )		
		Cry3Aa	NPTII	CP4 EPSPS
RBMT21-129	Leaf (n=18)	12.81 (9.12-16.53)	below LOD <sup>1</sup>	Not determined
	Tuber (n=12)	0.35 (0.11-0.90)	"	
RBMT21-350	Leaf (n=19)	20.54 (7.71-35.66)	"	Not determined
	Tuber (n=11)	0.28 (0.12-0.61)	"	
RBMT22-82	Leaf (n=18)	20.97 (14.97-29.44)	Not determined	28.34 (17.36-35.66)
	Tuber (n=12)	0.63 (0.49-0.79)		0.53 (0.21-0.78)

<sup>1</sup> limit of detection. The limit of detection was 0.3 ng/g tissue.

Using total protein levels of 1.6 and 2.0% for leaf and tuber fresh weight, respectively, the above protein expression levels correspond to the following % total protein.

**Table 5: Protein expression levels in tissue of Russet Burbank lines as a percentage of total protein**

Tissue	Protein expression levels (% total protein)		
	Cry3Aa	NPTII	CP4 EPSPS
Leaf	0.05-0.2	<0.0001	0.1-0.2
Tuber	0.0006-0.005	<0.0001	0.001-0.004

### **Conclusion**

The expression levels of all three novel proteins in potato tubers are low. Protein expression levels in tubers are significantly less than those in the corresponding leaf tissue.

### 3.4 *Impact on human health of the potential transfer of novel genetic material to cells of the human digestive tract*

The human health considerations in this regard depend on the nature of the novel genes and must be assessed on a case-by case basis.

In 1991, the World Health Organization (WHO) issued a report of a Joint FAO<sup>4</sup>/WHO Expert Consultation which looked at strategies for assessing the safety of foods produced by biotechnology (WHO 1991). It was concluded by that consultation that as DNA from all living organisms is structurally similar, the presence of transferred DNA in food products, in itself, poses no health risk to consumers.

The major concern in relation to the transfer of novel genetic material to cells in the human digestive tract is with antibiotic resistance genes. Antibiotic resistance genes can be present in some transgenic plants as a result of their use as marker genes to select transformed cells. It is generally accepted that there are no safety concerns with regard to the presence in the food of antibiotic resistance gene DNA *per se* (WHO 1993). There have been concerns expressed, however, that there could be horizontal gene transfer of antibiotic resistance genes from ingested food to microorganisms present in the human digestive tract and that this could compromise the therapeutic use of antibiotics.

This section of the report will therefore concentrate on evaluating the human health impact of the potential transfer of antibiotic resistance genes from New Leaf® Plus potatoes to microorganisms present in the human digestive tract.

Two antibiotic resistant genes have been transferred to the New Leaf® Plus potato lines — the *nptII* gene and the *aad* gene. The transferred *aad* gene may not be intact. The *nptII* gene confers resistance to the aminoglycoside antibiotics, neomycin, kanamycin, and geneticin (G418) and the *aad* gene confers resistance to the antibiotics spectinomycin and streptomycin. These antibiotics only have very limited clinical use. Neomycin is not used orally because of its toxicity but is still used topically in certain circumstances (Davis *et al* 1980). Streptomycin has mostly been replaced by newer aminoglycosides, although it is still used for special indications, such as in the treatment of tuberculosis and brucellosis (Kärenlampi 1996) and spectinomycin is rarely used clinically.

Lines RMBT21-129 and RMBT21-350 contain the *nptII* gene, under the control of the NOS promoter, meaning it will be expressed in plant cells but not bacterial cells, and line RMBT22-82 contains a copy or partial copy of the *aad* gene, under the control of a bacterial promoter.

The first issue that must be considered in relation to the presence of the *nptII* and *aad* genes in the New Leaf® Plus potatoes is the probability that these gene would be successfully transferred to and expressed in microorganisms present in the human digestive tract. The following steps are necessary for this to occur:

- excision of DNA fragments containing the antibiotic resistance gene;
- survival of DNA fragments containing the antibiotic resistance gene in the digestive tract;

---

<sup>4</sup> Food and Agriculture Organization.

- natural transformation of bacteria inhabiting the digestive tract;
- survival of the bacterial restriction system by the DNA fragment containing the antibiotic resistance gene;
- stable integration of the DNA fragment containing the antibiotic resistance gene into the bacterial chromosome or plasmid;
- maintenance and expression of antibiotic resistance gene by the bacteria. In the case of the *nptII* gene this would have to involve the acquisition of a bacterial promoter.

In the case of line RBMT22-82, there may be a slightly higher probability of horizontal gene transfer of the *aad* gene because of the transfer to the plant genome of a linked *Escherichia coli* origin of replication (*ori322*). Depending on the integrity of these components, the presence of these elements on the same DNA fragment could lead to the reconstitution of a plasmid capable of autonomous replication in *E. coli*. A plasmid is more likely to be successfully taken up than an isolated fragment of DNA. This however, would still be an extremely unlikely event.

The transfer of either the *nptII* or *aad* genes to microorganisms in the human digestive tract is therefore considered to be highly unlikely because of the number and complexity of the steps that would need to take place consecutively.

The second and most important issue that must be considered is the potential impact on human health in the unlikely event successful transfer of a functional antibiotic resistance gene to microorganisms in the human digestive tract did occur.

In the case of transfer of the *nptII* gene and the *aad* gene, the human health impacts are considered to be negligible. In the case of *nptII*, this gene occurs naturally in bacteria inhabiting the human digestive tract therefore the additive effect of an *nptII* gene entering the human gastrointestinal flora from a genetically modified plant would be insignificant compared to the population of kanamycin resistant microorganisms naturally present. In the case of the *aad* gene, this gene is common and can be found at high frequencies in natural populations of bacteria as well as clinical isolates (Shaw *et al* 1993). Natural populations of streptomycin resistant bacteria are far more likely to be sources of transferred antibiotic resistance than ingested plant material.

In relation to transfer of other novel genetic material to human cells via the digestive tract, this is also equally unlikely to occur. In considering the potential impact on human health, it is important to note that humans have always consumed large amounts of DNA as a normal component of food and there is no evidence that this consumption has had any adverse effect on human health. Furthermore, current scientific knowledge has not revealed any DNA sequences from ingested foods that have been incorporated into human DNA. Novel DNA sequences in genetically modified foods comprise only a minute fraction of the total DNA in the food (generally less than 0.01%) and are therefore unlikely to pose any special additional risks compared with the large amount of DNA naturally present in all foods.

## **Conclusion**

It is extremely unlikely that the kanamycin or streptomycin resistance genes would transfer from the New Leaf® Plus potatoes to bacteria in the human digestive tract because of the number and complexity of steps that would need to take place consecutively. In the highly unlikely event that the genes were transferred the human health impacts would be negligible because both antibiotic resistance genes are already commonly found in bacteria in the environment as well as inhabiting the human digestive tract and both antibiotics have very little, if any, clinical use in Australia and New Zealand.

It is also equally unlikely that other novel genetic material from the New Leaf® Plus potatoes would be transferred to human cells via the digestive tract. The novel genetic material in the potatoes comprises only a minute fraction of the total DNA therefore it is unlikely to pose any special additional risks compared with the large amount of DNA naturally present in all foods.

## **4. TOXICOLOGICAL ISSUES**

### **4.1 Levels of naturally-occurring toxins**

Wild tuberous *Solanum* species contain high concentrations of the toxic glycoalkaloids, which are very bitter in taste. The presence of glycoalkaloids in *Solanum* species is generally believed to be a natural plant defense mechanism against pests and diseases (Conner 1995). Modern potato cultivars accumulate high glycoalkaloid concentrations in green shoot tissue and in tubers upon exposure to light. In some cultivars, significant concentrations of glycoalkaloids can also accumulate in tubers not exposed to light. The variation in glycoalkaloid content of tubers can be attributed to both genetic effects and the environmental conditions under which the plants are grown and stored following harvest (van Gelder 1990). The concentration of glycoalkaloids in potato tubers in advanced lines of modern breeding programs is usually routinely monitored (Morris and Lee 1984).

Studies submitted by Monsanto:

Lavrik, P.B. *et al* (1997). Compositional analysis of potato tubers derived from Colorado potato beetle and potato leafroll virus resistant (New Leaf® Plus) Russet Burbank potato plants grown under field conditions. Monsanto Study No. 96-01-37-23.

The New Leaf® Plus potato lines were grown in six locations across the USA during 1995 and 1996. Tubers were collected at harvest from four replicated plots from each site and analysed for glycoalkaloid levels. The controls were tubers isolated from non-transformed Russet Burbank potato lines grown at the same field location. The field trials were carried out in a randomised complete block design consisting of eight to fifteen replicated plots per line. This study was limited to tubers collected from four replicates per site. A summary of the results is presented in Table 6 below.

**Table 6: Total glycoalkaloid content<sup>#</sup> of tubers from New Leaf® Plus and control lines.**

Potato line	Total glycoalkaloids (mg/100 g fresh weight)
RBMT21-129	5.4 (2.1-8.9)
RBMT21-350	4.8 (1.7-16.3)
RBMT22-82	5.1 (2.8-17.5)
Russet Burbank control	4.3 (2.6-25.0)
Literature range (Russet Burbank)	3.1-16.1

<sup>#</sup> mean values, range in brackets (n=24, except for RBMT21-129 where n=20)

The data presented in Table 6 demonstrates that the New Leaf® Plus potato lines have total glycoalkaloid levels which are slightly elevated compared to those of the non-transformed Russet Burbank control line. These differences, however, are not statistically significant. The range of levels for glycoalkaloids reported for both the New Leaf® Plus and control lines varied widely but are generally comparable to the literature reported range for Russet Burbank cultivars. There are no safety concerns regarding the slightly elevated glycoalkaloid levels in the New Leaf® Plus lines.

#### 4.2 Potential toxicity of novel proteins

All three New Leaf® Plus potato lines express the Cry3Aa protein and the data suggests, although there is no direct evidence, that they may also express the PLRV replicase. In addition to these two proteins, lines RBMT21-129 and RBMT21-350 also express NPTII and line RBMT22-82 expresses CP4 EPSPS. This section of the report will therefore assess the potential toxicity of all four proteins.

Studies submitted by Monsanto<sup>#</sup>:

Lavrik, P.B. and Grace, A.M. (1997). Assessment of the equivalence of *B.t.t.*, NPTII and CP4 EPSPS proteins produced in *E. coli* and in Russet Burbank potato plant lines resistant to potato leafroll virus and Colorado potato beetle. Monsanto Study No. 96-01-37-26.

Berberich, S.A. *et al* (1993). Preparation and verification of dose for a mouse acute oral toxicity study with neomycin phosphotransferase II protein (NPTII). Monsanto Study ML-91-409.

Reding, H.K. (1998). Safety of PLRV replicase in NewLeaf® Plus potatoes. Monsanto Company. EPA application No. 198176.

Naumovich, L. (1994) The infection of the *Solanum tuberosum*, Russet Burbank potato, by PVX, PVY, and PLRV viruses during the cultivation of the tuber. A biology study presented to the Monsanto/St Louis Post-Dispatch Greater St Louis Science Fair.

<sup>#</sup>Refer also to studies submitted for Applications A338 – Roundup Ready soybeans (CP4 EPSPS) and A382 – New Leaf® potatoes (Cry3Aa, NPTII)

#### Cry3Aa

Cry3Aa is insecticidal only to Coleopteran insects (MacIntosh *et al* 1990) and its specificity of action is directly attributable to the presence of specific receptors in the target insects (Wolfersberger 1990, Ferré *et al* 1991). There are no receptors for the  $\delta$ -endotoxins of *B. thuringiensis*, including Cry3Aa, on the surface of mammalian intestinal cells (Hubert *et al* 1995), therefore, humans, as well as other mammals, are not susceptible to this protein.

The potential toxicity of Cry3Aa was assessed for New Leaf® potatoes under Application A382 where acute oral toxicity studies in mice were submitted for evaluation. These studies are also relevant to this application as the gene construct for the *cry3Aa* gene used in the New Leaf® Plus potatoes will give rise to an identical protein to that produced in the New Leaf® potatoes. Acute tests are used because it is known that protein toxins generally act via acute mechanisms (Jones and Maryanski 1991). For a detailed summary of the acute toxicity study, refer to the safety evaluation for Application A382 – New Leaf® potatoes. A brief summary of the findings is presented below.

The Cry3Aa protein used in the toxicity study was produced in *E. coli* because the plant lines did not express enough protein for purification of large quantities for toxicity testing. Data was presented to indicate that the bacterially produced Cry3Aa is equivalent to the plant produced Cry3Aa in terms of its molecular mass, N-terminal amino acid sequence, lack of glycosylation, and biological activity. Therefore, the *E. coli* produced Cry3Aa is considered to be a suitable substitute for plant produced Cry3Aa.

The Cry3Aa protein was administered by gavage to CD-1 mice at doses up to 5220 mg/kg body weight, the mice being observed for a period of seven days. No abnormal clinical signs were observed in the mice during the study that could be attributed to the treatment. No significant differences were observed in body weight, cumulative body weight or food consumption. Several minor pathologic changes were observed at necropsy but these were randomly distributed among all groups and could not be attributed to the treatment. On the basis of these findings the Cry3Aa protein was considered to be non-toxic.

#### **PLRV replicase protein**

Expression studies using Western blot analysis were not able to detect the presence of the PLRV replicase in the New Leaf® Plus potatoes. Other evidence (discussed in Section 3.3 above), however, suggests that the replicase protein is expressed. In the light of only equivocal evidence for non-expression of the PLRV replicase, it is necessary to also consider the potential toxicity of the PLRV replicase.

The applicant did not submit any data from acute oral toxicity studies with the PLRV replicase protein. Instead, the applicant has provided the following arguments and evidence for the safety of the PLRV replicase protein.

#### *History of safe consumption*

PLRV must produce its replicase protein to be able to replicate in potatoes. Although it has not been possible to detect viral replicase in PLRV-infected potatoes, it logically follows that the replicase must be present in plants where the virus can be detected. PLRV infection of potatoes has been known to occur since the early twentieth century (Slack 1993). In some potato crops in the United States, the incidence of PLRV infection routinely approaches 100% if insecticides to control the aphid vector of the virus are not used (Thomas 1997). In a survey undertaken on behalf of the applicant, between 7-14% of fresh potatoes in supermarkets in the United States were found to be infected with PLRV. Therefore, it can be said that there is a long history of human exposure to the PLRV replicase protein through the consumption of PLRV-infected potatoes, without any reported adverse effects.

### *PLRV replicase levels in New Leaf® Plus potatoes*

The applicant provided evidence which suggests that the level of replicase protein expressed in the New Leaf® Plus potatoes is likely to be lower than what would be expressed in PLRV-infected potatoes. This comes from data that New Leaf® Plus potatoes produce <0.05ng *PLRVrep* mRNA/mg total RNA compared to the 3.4ng viral RNA/mg total RNA produced by PLRV-infected plants. It can be inferred from this data that the New Leaf® Plus potatoes will contain substantially lower amounts of PLRV replicase than that present in PLRV-infected potatoes.

### *Similarity with known toxins*

The amino acid sequence of the PLRV replicase was compared to a database of 2625 known toxins. No significant amino acid similarity was observed.

### *Conclusion*

There is a long history of safe human exposure to the PLRV replicase protein through the consumption of PLRV-infected potatoes. In addition, the evidence suggests that exposure to the PLRV replicase protein from the consumption of the New Leaf® Plus potatoes is likely to be lower than that from consumption of PLRV-infected potatoes. Furthermore, there is no significant similarity between the PLRV replicase and any known protein toxins. On the basis of this evidence, animal studies of the toxicity of the PLRV replicase protein are considered to be unnecessary and it is concluded that the PLRV replicase protein, as expressed in the New Leaf® Plus potatoes, is non-toxic to humans.

### **Neomycin phosphotransferase II**

The potential toxicity of NPTII was assessed for New Leaf® potatoes under Application A382 where acute oral toxicity studies in mice were submitted for evaluation. These studies are also relevant to this application as the gene construct for the *nptII* gene used in the New Leaf® Plus potatoes will give rise to an identical protein to that produced in the New Leaf® potatoes. For a detailed summary of the toxicity study, refer to the safety evaluation for Application A382 – New Leaf® potatoes. A brief summary of the findings is presented below.

The NPTII protein used in the study was produced from *E. coli* because the plant lines did not express enough protein for purification of large quantities for toxicity testing. Data was presented to show that the *E. coli* produced NPTII is equivalent to the plant produced NPTII in terms of its molecular mass, N-terminal amino acid sequence, lack of glycosylation, and biological activity. The *E. coli* produced NPTII is therefore considered to represent a suitable substitute for plant produced NPTII.

The NPTII protein was administered by gavage to mice at doses up to 5000 mg/kg body weight for a period of 8-9 days. There were no statistically significant differences in mean body weights or cumulative body weight gain in any of the treated groups. No abnormal clinical signs were noted, there were no unscheduled deaths and there were no differences in mean terminal body weights. No gross lesions were observed at necropsy. On the basis of these findings NPTII was considered to have low oral toxicity.

## **CP4 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS)**

The *CP4 EPSPS* gene construct used for the production of the New Leaf® Plus potato lines encodes a protein that is identical to the CP4 EPSPS protein produced in Roundup Ready soybeans. The potential toxicity of CP4 EPSPS was previously assessed for Roundup Ready soybeans under Application A338 where it was concluded to be of low toxic potential.

For Roundup Ready soybeans, the potential toxicity of CP4 EPSPS was assessed by three methods — firstly, through an acute oral toxicity study in mice, secondly through comparison of its amino acid sequence with the amino acid sequences of known toxins and lastly by its similarity with other EPSPS proteins which have a safe history of human consumption. The findings of these evaluations are summarised briefly below.

### *Acute oral toxicity study*

The CP4 EPSPS protein, lacking the chloroplast transit peptide, was administered by gavage to mice at doses up to 572 mg/kg body weight. No adverse effects were observed at these dose levels.

### *Comparison with known toxins*

CP4 EPSPS was compared to the amino acid sequence of 1935 known protein toxins. No meaningful similarity was found other than would be expected given that certain functional domains are generally conserved between proteins.

### *Similarity to other EPSPS proteins*

All crop, fungal and microbial food sources such as Baker's yeast (*Saccharomyces cerevisiae*) or *Bacillus subtilis* contain EPSPS proteins (Kishore *et al* 1988) and, as such, EPSPS proteins have a long history of safe human consumption. The CP4 EPSPS protein shares significant amino acid sequence homology with other EPSPS proteins therefore it is unlikely to produce any toxic effects in humans.

### *Conclusion*

As the CP4 EPSPS protein expressed in New Leaf® Plus potato line RBMT22-82 is identical to the EPSPS protein expressed in Roundup Ready soybeans, the conclusions of that safety evaluation are relevant here. No additional data or evidence has emerged which would necessitate revising the conclusion of the previous study. Furthermore, the CP4 EPSPS protein has itself now been consumed for at least five years in Roundup Ready soybeans without any reports of adverse effects. Therefore, it can be concluded that the CP4 EPSPS protein is non-toxic to humans.

## **4.3 Levels of naturally-occurring allergenic proteins**

Potatoes are not generally regarded as major sources of food allergy, although patatin, the main storage protein of potatoes, has recently been reported to induce an allergic reaction in some individuals (Seppälä *et al.*, 1999). The clinical importance of patatin as a food allergen has yet to be confirmed.

As potatoes are not classified as major sources of food allergy, and there have yet to be any confirmed potato allergens described, an assessment of the naturally-occurring allergenic proteins of New Leaf® Plus potatoes is unnecessary.

#### **4.4 Potential allergenicity of novel protein**

The concerns regarding potential allergenicity of novel proteins are two fold. Firstly, there are concerns that the ability to express new or different proteins in food will result in the transfer of allergens from one food to another, thereby causing some individuals to develop allergic reactions to food they have not previously been allergic to. Secondly, there are concerns that the transfer of novel proteins to food will lead to the development of new allergies in certain individuals. The former is more easily addressed than the latter because if an allergen is already known it is possible, using human sera or human skin tests, to test if it has been transferred. There are no reliable tests or animal models, however, which enable the prediction of the allergenic potential of novel proteins. Instead, potential allergenicity can only be indicated by examination of a number of characteristics of the novel protein, such as whether it is derived from a known allergenic source, its physical/chemical characteristics (most allergens have a molecular mass of between 10 and 70 kDa, are glycosylated and are resistant to acid and protease degradation), whether it has any sequence similarity to any known allergens, and whether it is likely to be present in large amounts in the food as consumed and therefore have potential for allergic sensitisation.

There are four new proteins expressed in the New Leaf® Plus potatoes – Cry3Aa, PLRV replicase, NPTII, and CP4 EPSPS. The potential allergenicity of Cry3Aa and NPTII was considered for New Leaf® potatoes, under Application A382 and the potential allergenicity of the CP4 EPSPS protein was considered for Roundup Ready soybeans under Application A338. The findings of those assessments are briefly summarised.

##### **Cry3Aa, NPTII and CP4 EPSPS**

For Cry3Aa, NPTII and CP4 EPSPS it was concluded that each protein is within the size range of known allergens, however, none are glycosylated and all are rapidly degraded in the proteolytic and acid conditions of simulated gastric fluid suggesting they would not survive mammalian digestion. None of the proteins have any significant similarity to known allergens, nor are they present in large amounts in potato tubers. On the basis of this data and on the basis that humans have a prior history of exposure to these proteins with no recorded instances of allergenicity, it was concluded that Cry3Aa, NPTII and CP4 EPSPS are unlikely to be allergenic to humans. No additional data or evidence has emerged which would necessitate revising this conclusion.

##### **PLRV replicase**

The potential allergenicity of the PLRV replicase protein has not previously been considered.

##### *Source of the protein*

The PLRV replicase is encoded by the *PLRVrep* gene, which was derived from a PLRV isolate obtained from a naturally infected potato in the United States (Kaniewski *et al* 1995). The PLRV replicase protein in the New Leaf® Plus potatoes is therefore identical to that which occurs in PLRV infected potatoes. PLRV infection of potatoes is a common

occurrence (see Section 4.2). Infected potatoes are still suitable for human consumption. PLRV replicates in the vascular tissues of the tuber (Weidemann and Casper 1992) therefore both the virus and its expressed proteins, including the replicase, are routinely consumed. Despite the widespread consumption of PLRV-infected tubers, there have been no reports of allergic reactions to any of the viral proteins.

#### *Physical and chemical characteristics*

The PLRV replicase protein has a molecular weight of 130 kDa, which is outside the size range of known allergens (10-70 kDa). However, it is speculated that the replicase protein undergoes post-translation processing, therefore the processed products could well fall within the size range of known allergens. The PLRV replicase, however, is not glycosylated as it is expressed in the plant cell cytoplasm. For glycosylation of the PLRV replicase to occur, the protein would need to be transported through the endoplasmic reticulum and Golgi bodies (Taiz and Zeiger 1991). This requires the presence of specific targeting sequences on the protein and none of these were included in the *PLRVrep* gene construct.

#### *Similarity to known allergens*

The amino acid sequence of the PLRV replicase protein was compared to a database of 265 known allergen and gliadin sequences. In agreement with current understanding of allergen epitope structure (O’Hehir *et al* 1991), a significant sequence similarity was defined as a sequence identity of greater than seven contiguous amino acids. No significant sequence identity with any of the 265 allergens was found.

#### *Presence of the protein in food as consumed*

One of the factors contributing to the allergenicity of certain proteins is their high concentration in foods that elicit an allergic response (Taylor *et al* 1987, Taylor 1992, Taylor *et al* 1992). This is true for milk (Baldo, 1984, Taylor *et al* 1987), soybean (Shibasaki *et al* 1980, Burks *et al* 1988, Pendersen and Djurtoft 1989) and peanuts (Barnett *et al* 1983, Sachs *et al* 1981, Barnett and Howden 1986, Kemp *et al* 1985).

The PLRV replicase protein, in contrast, cannot be detected in tubers of New Leaf® Plus potatoes. There is only equivocal evidence that the protein is expressed at all.

### **Conclusion**

It has previously been concluded that Cry3Aa, NPTII and CP4 EPSPS are unlikely to be allergenic to humans. No additional data or evidence has emerged which would necessitate revising this conclusion.

The PLRV replicase protein in the New Leaf® Plus potatoes has no significant similarity to any known allergens, nor is it present in large quantities in potato tubers. On the basis of this evidence and the fact that humans have a long history of exposure to this protein, through the consumption of PLRV-infected potatoes, with no recorded instances of allergenicity, it can be concluded that the PLRV replicase protein is unlikely to be allergenic to humans.

## 5. NUTRITIONAL ISSUES

### 5.1 *Nutrient analysis*

There are concerns that genetic modification will affect the overall nutritional composition of a food, or cause unintended changes that could adversely affect the safety of the product. Therefore a safety assessment of food produced from genetically modified plants must include analysis of the composition of the food, based on a comparison with other commercial varieties of the crop. Generally, comparisons are made not only with the parental line but also with other non-transformed lines. If the parameter for the transformed line is within the normal range for non-transformed lines, this is considered acceptable (Hammond and Fuchs 1998).

In undertaking a compositional analysis of potatoes there are a number of key defining nutrients and constituents that should be measured as part of that analysis. They are total tuber solids (measured as tuber dry matter), sugars, protein and vitamin C. Tuber solids are an important quality factor for processing and are also the single most important determinant of culinary appeal (Murphy *et al* 1967). Approximately 75% of the dry matter content of potatoes consists of starch. The remainder is composed of sugars, protein, and assorted cell and cell wall components (Storey and Davies, 1992). The major sugars in potatoes are sucrose as well as the reducing sugars fructose and glucose. They are present in small quantities and are inconsequential as a source of energy. However, like total solids, they are a very important factor in processed food quality. Potatoes also contain measurable amounts of proteins, fats, carbohydrates, and numerous vitamins and minerals. However, they are only a significant dietary source for two of these constituents – protein and vitamin C (Storey and Davies 1992, Pennington and Wilkening 1997). Potato proteins are highly digestible, have a fairly good balance of amino acids and are especially high in the essential amino acid lysine. Measurement of total protein is considered more informative than measurement of individual amino acids as nearly all of the proteins in potato tubers (albumin, globulin, glutelin, and prolamin) have a similar amino acid composition, therefore, changes in their respective ratios will have little impact on the amino acid profile (Storey and Davies, 1992).

Studies submitted by Monsanto:

Lavrik, P.B. (1995). Compositional analysis of potato tubers derived from Colorado potato beetle and potato leafroll virus resistant (NewLeaf®) Russet Burbank potato plants grown under field conditions. Monsanto Study No. 96-01-37-23.

Rogan, G. J. *et al.* (1999). Composition analysis of potato tubers from NewLeaf® Y and NewLeaf® Plus potato plants grown under field conditions. Monsanto Study No. 98-01-37-27.

The applicant undertook two separate field studies of the New Leaf® Plus potatoes. The first study was conducted in 1995 and 1996 at three locations in the United States and three locations in Canada. At each location, eight to fifteen replicated plots were grown per line. Tubers were collected at harvest from four replicated plots and analysed for total solids, dextrose, sucrose, vitamin C, soluble protein, and proximate composition (total protein, fat, crude fibre, ash, total carbohydrates and calories). The controls were tubers isolated from non-transformed Russet Burbank potato lines grown at the same field location. The field trials were carried out in a randomised complete block design consisting of eight to fifteen replicated plots per line. This study was limited to tubers collected from four replicates per site.

The second study was conducted in 1998 at three locations in the United States. Four replicated plots were grown at one of the sites, whereas plants were grown in non-replicated plots at the other two sites. Compositional analyses were done of amino acid, vitamin B<sub>6</sub>, niacin, copper, magnesium, and potassium content.

### Major constituents

Summaries of the results of proximate and other major constituent analyses are presented in Table 7 and 8 below.

**Table 7: Proximate and major constituent content<sup>1</sup> of New Leaf® Plus potatoes – Canadian data**

Constituent	New Leaf® Plus lines			RB Control	Lit. range
	RBMT21-129	RBMT21-350	RBMT22-82		
<b>Total solids (% fw)</b>	21.1 (20.4-22.4)	21.6 (20.2-23.7)	21.0 (20.2-22.5)	21.7 (19.9-23.5)	16.8-24.5
<b>Sugars (%fw):</b>					
Dextrose	0.24 (0.05-0.35)	0.21 (0.05-0.38)	0.24 (0.08-0.37)	0.24 (0.05-0.40)	0.04-0.52
Sucrose	0.14 (0.10-0.20)	0.14 (0.11-0.18)	0.14 (0.11-0.16)	0.16 (0.12-0.22)	0.1-0.88
<b>Soluble protein (% dw)</b>	4.6 (4.1-5.4)	4.6 (4.0-5.2)	4.5 (3.9-5.1)	4.4 (3.9-4.7)	3.4-7.3
<b>Proximate<sup>2</sup>:</b>					
Moisture	1.39 (1.03-1.42)	1.43 (1.02-1.68)	1.20 (1.01-1.44)	1.48 (1.03-2.71)	-
Total protein	9.86 (7.16-12.75)	9.94 (7.6-13.4)	9.93 (8.20-11.98)	9.90 (7.81-12.59)	7.1-14.6
Fat	0.20 (0.13-0.32)	0.19 (0.10-0.30)	0.20 (0.13-0.27)	0.16 (0.11-0.23)	0.1-0.8
Ash	4.68 (4.21-4.85)	4.70 (4.08-5.72)	4.78 (3.95-5.69)	4.75 (3.93-5.43)	2.2-9.5
Crude fibre	1.64 (1.15-1.78)	1.55 (1.12-1.78)	1.61 (1.24-1.79)	1.68 (1.39-2.15)	0.2-3.5
Carbohydrates	85.2 (81.2-86.5)	85.2 (82.5-86.6)	85.1 (81.7-85.7)	85.2 (80.74-86.84)	84.5 (avg)
Calories	378 (376-379)	377 (372-380)	377 (374-381)	376 (370-380)	350 (avg)

<sup>1</sup> mean values, range in brackets (n=12 except for RBMT21-129 where n=8)

<sup>2</sup> except for moisture and calories, reported values are in g/100 g dry weight. Moisture is reported in g/100 g of lyophilised tuber powder. Calories are reported in calories/100 g dry weight.

**Table 8: Major constituents<sup>#</sup> of New Leaf® Plus potatoes – USA data**

Constituent	New Leaf® Plus lines			RB Control	Lit. range
	129	350	82		
<b>Total solids (% FW)</b>	21.6 (20.3-23.0)	21.9 (20.6-24.1)	21.0 (19.4-22.1)	21.5 (20.5-22.5)	16.8-24.5
<b>Sugars (%FW):</b>					
Dextrose	0.09 (0.04-0.12)	0.09 (0.06-0.16)	0.11 (0.05-0.16)	0.10 (0.07-0.13)	0.04-0.52
Sucrose	0.18 (0.14-0.23)	0.20 (0.16-0.25)	0.18 (0.13-0.23)	0.20 (0.14-0.26)	0.1-0.88
<b>Soluble protein (% DW)</b>	5.0 (4.4-5.8)	5.1 (4.6-5.5)	5.0 (4.6-5.8)	5.0 (4.5-5.6)	3.4-7.3

<sup>#</sup> mean values, range in brackets (n=12, except for RBMT21-350 where n=11)

There were no statistically significant differences between New Leaf® Plus potatoes and the control lines for any of the major constituents and the levels reported were all comparable to the normal ranges for Russet Burbank cultivars.

### Amino acid content

The concentration of 18 out of a total of 20 amino acids was measured for the New Leaf® Plus potato lines. The two amino acids not analysed were asparagine and glutamine. The data obtained on the amino acid composition of the New Leaf® Plus potato lines is summarised in Table 9 below.

**Table 9: Amino acid content<sup>#</sup> of New Leaf® Plus Russet Burbank potato lines**

Amino acid	New Leaf® Plus lines			Control	Lit. range
	RBMT21-129	RBMT21-350	RBMT22-82		
	(mg/200 g tuber fresh weight)				
Aspartic acid	1280 (812-1822)	1096 (681-1419)	1296 (815-1982)	1250 (728-1630)	677-1476
Threonine	147 (116-193)	134 (106-162)	152 (115-211)	147 (119-173)	102-214
Serine	151 (109-197)	147 (121-175)	159 (121-215)	155 (124-185)	125-255
Glutamic acid	771 (550-1017)	701 (491-858)	779 (588-1115)	793 (516-1055)	583-1207
Proline	124 (92-160)	111 (78-141)	129 (84-186)	119 (88-160)	89-366
Glycine	122 (101-148)	112 (96-127)	126 (102-178)	121 (107-143)	92-195
Alanine	118 (96-147)	112 (92-133)	124 (101-171)	117 (99-135)	87-238
Cysteine	62 (56-67)	59 (55-63)	64 (56-78)	62 (57-70)	48-93
Valine	217 (171-248)	199 (162-245)	237 (192-374)	218 (175-284)	196-363
Methionine	59 (48-66)	52 (42-62)	62 (48-100)	56 (41-84)	57-100
Isoleucine	140 (109-162)	123 (100-151)	148 (117-230)	139 (117-178)	119-238
Leucine	224 (173-286)	200 (156-240)	229 (169-323)	220 (176-263)	171-346
Tyrosine	128 (97-162)	129 (111-150)	149 (122-209)	144 (117-178)	114-236
Phenylalanine	167 (134-203)	156 (130-184)	180 (141-268)	168 (133-208)	138-272
Histidine	78 (64-95)	72 (57-83)	84 (67-123)	82 (66-100)	33-117
Lysine	235 (194-284)	211 (175-241)	245 (198-360)	233 (193-291)	154-342
Arginine	196 (147-253)	180 (131-219)	215 (170-320)	200 (145-254)	175-362
Tryptophan	43 (38-46)	39 (36-42)	45 (37-64)	42 (34-54)	29-70

<sup>#</sup> mean values, range in brackets (n=6).

The levels reported for amino acids in the New Leaf® Plus potatoes are comparable to the control values as well as the literature reported ranges.

### Vitamin and mineral content

Data obtained on the vitamin and mineral composition of the New Leaf® Plus potato lines is summarised in Table 10 below.

**Table 10: Mean vitamin and mineral content<sup>#</sup> of New Leaf® Plus potato lines**

Vitamin/ mineral <sup>1</sup>	New Leaf® Plus lines			RB Control	Lit. range
	RBMT21-129	RBMT21-350	RBMT22-82		
<b>Vitamins:</b>					
Vitamin C (Can)	16.7 (14.2-17.9)	16.8 (14.3-20.2)	16.8 (15.3-19.3)	16.4 (14.2-18.6)	10.3-22.0
Vitamin C (USA)	10.1 (8.2-13.0)	9.9 (8.2-11.3)	10.4 (7.6-12.8)	10.0 (8.8-12.6)	"
Vitamin B6	0.51 (0.32-0.58)	0.56 (0.31-0.57)	0.52 (0.30-0.78)	0.52 (0.45-0.56)	0.26-0.82
Niacin	4.03 (2.81-5.10)	3.99 (3.20-4.44)	4.28 (2.67-5.11)	4.06 (3.49-4.60)	0.18-6.2
<b>Minerals:</b>					
Copper	0.39 (0.14-0.61)	0.30 (0.14-0.50)	0.33 (0.11-0.64)	0.32 (0.14-0.50)	0.03-1.4
Magnesium	52.2 (48.5-56.6)	45.7 (43.0-50.8)	46.9 (38.5-67.8)	61.5 (47.1-66.1)	22.5-110
Potassium	1072.6 (955.9-1185.6)	1026.6 (966.7-1120.2)	1038.2 (931.2-1512.6)	1080.7 (979.2-1202.7)	700-1250

<sup>1</sup> Except for vitamin C content, values are reported as mg/200 g fresh weight. Vitamin C content is reported as mg/100 g fresh weight.

<sup>#</sup> mean values, range in brackets (n=6, except for vitamin C content where the number of samples corresponds to those for tables 5 and 6)

There were no statistically significant differences between New Leaf® Plus potatoes and the control line for any of vitamins or minerals analysed and the levels reported were comparable to the literature reported ranges for Russet Burbank cultivars.

## **Conclusion**

Based on the data submitted in the present application, The New Leaf® Plus potato lines are compositionally equivalent to other commercial Russet Burbank potato varieties.

### **5.2 *Levels of anti-nutrients***

The only known anti-nutrient present in potato is trypsin inhibitor. Trypsin inhibitors are classed as anti-nutrients because they interfere with the digestion of proteins leading to decreased animal growth. Trypsin inhibitors are heat labile and are destroyed during the cooking process or during processing when heat treatment is applied.

As heating inactivates trypsin inhibitor, its presence is only an issue when raw potatoes are consumed. Humans rarely consume raw potatoes due to the indigestibility of the ungelatinised starch.

### **5.3 *Ability to support typical growth and well-being***

In assessing the safety of food produced using gene technology, a key factor is the need to establish that the food is nutritionally adequate and will support typical growth and well-being. In most cases, this can be achieved through an understanding of the genetic modification and its consequences together with an extensive compositional analysis of the food. Where, on the basis of available data, there is still concern or doubt in this regard, carefully designed feeding studies in animals may provide further re-assurance that the food is nutritionally adequate. Such studies may be considered necessary where the compositional analysis indicates significant differences in a number of important components or nutrients or where there is concern that the bioavailability of key nutrients may be compromised by the nature of the genetic changes to the food.

The compositional and other data presented in the application are considered adequate for establishing the ability of New Leaf® Plus potatoes to support typical growth and well-being. Additional studies, including animal feeding studies, are therefore not required.

## **6. OTHER ISSUES**

### **6.1 *Estimation of dietary intake of novel proteins***

If the concentration of a substance in a food is known and data is available on the human consumption of that food then it is possible to estimate the dietary intake of that substance for the population. In safety assessments, dietary intakes are usually only estimated in circumstances where a substance is considered to be hazardous. In this way it is possible to determine the likely human exposure to the hazard and thus ascertain whether there is cause for concern.

None of the novel proteins in the New Leaf® Plus potatoes are considered to be hazardous therefore a dietary exposure assessment is unnecessary for determining if there is cause for concern. However, such information can provide additional assurance that exposure to the novel protein is low and/or that the novel protein is likely to be present in the diet at levels well below those found to be safe in animal toxicity studies.

The concentration of Cry3Aa and CP4 EPSPS in the New Leaf® Plus potatoes is known but the concentration of the NPTII and PLRV replicase proteins was unable to be quantified, therefore it is possible to only estimate the dietary intake for Cry3Aa and CP4 EPSPS.

Cry3Aa is expressed in the New Leaf® Plus potato tubers at levels ranging from 0.11 to 0.9 µg protein/g fresh weight and CP4 EPSPS is expressed at levels ranging from 0.21 to 0.78 µg protein/g fresh weight (see Table 2, Section 3.3).

Australian and New Zealand consumption data is available for potato crisps, instant mashed potato, and potato fries, although no data is currently available for potato flour and potato starch. The consumption data is presented in Table 11 below.

**Table 11: Estimated consumption of processed potato products in Australia and New Zealand.**

Food	Country	All respondents (g/day)		Consumers only (g/day)		
		mean		mean	median	95 <sup>th</sup> percentile
Potato crisps	Aus	2.8		38.8	25	100
	NZ	2.9		48.4	40	150
Instant mashed potato	Aus	-		-	-	-
	NZ	0.007		34.6	34.6	34.6
Potato fries, commercial	Aus	16.6		132.5	113	264
	NZ	18.6		141.2	142	300
Total potato products	Aus	19.4		-	-	-
	NZ	21.5		118	112.2	300

For calculation of the dietary intake of the novel proteins, the highest potato consumption figure (300 g/day) and the highest protein concentration was used. This represents a ‘worst case’ estimate and also makes allowances for the lack of consumption data for potato flour and potato starch.

To do the calculation, assumptions about the proportion of processed potato products derived from the New Leaf® Plus potatoes must be made. Data on market penetration of the New Leaf® Plus potatoes is not available. In the absence of information about market penetration, two estimates are made — one using a very worst case estimate where it is assumed that all potato products are derived entirely from New Leaf® Plus potatoes and the other, probably more realistic estimate, where it is assumed that 10% of potato products are derived from New Leaf® Plus potatoes. The two estimates of dietary intake for Cry3Aa and CP4 EPSPS are presented in Table 12 below.

**Table 12: Estimate of dietary intake of Cry3Aa and CP4 EPSPS**

Novel protein	Estimated dietary intake			
	100 % market penetration		10 % market penetration	
	µg /day	µg/kg BW/day <sup>1</sup>	µg /day	µg/kg BW/day
Cry3Aa (0.11-0.9µg/g FW)	33-270	0.51-4.2	3.3-27.0	0.05-0.42
CP4 EPSPS (0.21-0.78 µg/g FW)	63-234	0.97-3.6	6.3-23.4	0.096-0.36

<sup>1</sup> assuming a body weight of 65 kg.

For Cry3Aa, the very worst-case estimate is at least 1.2 million times less than the dose found to have no adverse effects in mice (5220 mg Cry3Aa/kg BW, administered as two doses in a single day). For CP4 EPSPS, the estimate is at least 150 000 times less than the dose found to have no adverse effects in mice (572 mg CP4 EPSPS/kg BW, administered as a single dose). Therefore, even if all processed potato products were to be derived from the New Leaf® Plus potatoes, a very large margin of safety exists for both proteins.

## **ACKNOWLEDGEMENTS**

ANZFA gratefully acknowledges the expert comments on the safety assessment of food derived from insect and PLRV-protected potato lines RBMT21-129, RBMT21-350 and RBMT22-82 provided by Associate Professor Richard T. Roush, Centre for Weed Management Systems, Waite Institute, University of Adelaide, Glen Osmond 5064, and Professor Tony Connor, New Zealand Institute for Crop and Food Research Ltd and Soil, Plant and Ecological Sciences Division, Lincoln University, New Zealand.

## REFERENCES

- Almeida, E.R.P., Gossele, V., Muller, C.G., Dockx, J., Reynaerts, A., Botterman, J., Krebbers, E. and Timko, M.P. (1989). Transgenic expression of two marker genes under the control of an *Arabidopsis rbcS* promoter: sequences encoding the Rubisco transit peptide increase expression level. *Mol. Gen. Genet.* **218**: 78-86.
- ANZFA (1999). Guidelines for the safety assessment of foods to be included in Standard A18 — food produced using gene technology.
- Bachmann, B.G. (1987). Derivatives and genotypes of some mutant derivatives of *Escherichia coli* K-12. In: *Escherichia coli and Salmonella typhimurium*, Neidhardt, F.C. (ed), Volume 2, American Society for Microbiology, Washington, DC, pp 1190-1219.
- Baldo, B.A. (1984). Milk allergies. *Austral. J. Dairy Technolog.* **39**: 120-128.
- Barker, R.F., Idler, K.B., Thompson, D.V. and Kemp, J.D. (1983). Nucleotide sequence of the T-DNA region from *Agrobacterium tumefaciens* octopine Ti plasmid pTi5955. *Plant Mol. Biol.* **2**: 335-350.
- Barnett, D., Baldo, B.A. and Howden, M.E.H. (1983). Multiplicity of allergens in peanuts. *J. Allergy Clin. Immunol.* **72**: 61-68.
- Barnett, D. and Howden, M.E.H. (1986). Partial characterisation of an allergenic glycoprotein from peanut (*Arachis hypogaea* L.). *Biochem. Biophys. Acta* **882**: 97-105.
- Barry, G., Kishore, G., Padgett, S., Taylor, M., Kolacz, K., Weldon, M., Eichholtz, Re. D., Fincher, K. and Hallas, L. (1992). Inhibitors of amino acid biosynthesis: strategies for imparting glyphosate tolerance to crop plants. In: *Biosynthesis and Molecular Regulation of Amino Acids in Plants*, Singh, B.K., Flores, H.E. and Shannon, J.C. (eds), American Society of Plant Physiologists, Rockville, Maryland, pp139-145.
- Bazan, J.F. and Fletterick, R.J. (1989). Comparative analysis of viral cysteine protease structural models. *FEBS Lett.* **249**: 5-7.
- Beck, E., Ludwig, G., Auerswald, E.A., Reiss, B. and Schaller, H. (1982). Nucleotide sequence and exact localisation of the neomycin phosphotransferase gene from transposon Tn5. *Gene* **19**: 327-336.
- Bevan, M., Barnes, W.M. and Chilton, M. (1983). Structure and transcription of the nopaline synthase gene region of T-DNA. *Nucleic Acids Res.* **11**: 369-385.
- Bolivar, F., Rodriguez, R.L., Greene, P.J., Betlach, M.C., Heyneker, H.L. and Boyer, H.W. (1977). Construction and characterisation of new cloning vehicles II. A multipurpose cloning system. *Gene* **2**: 95-113.
- Burks, A.W., Jr., Brooks, J.R. and Sampson, H.A. (1988). Allergenicity of major component proteins of soybean determined by enzyme-linked immunosorbent assay (ELISA) and immunoblotting in children with atopic dermatitis and positive soy challenges. *J. Allergy Clin. Immunol.* **81**: 1135-1142.

- Burton, W.G. (ed) (1987) *The Potato*. Langman Sci. and Technical, New York. 742pp.
- Cole, D.J. (1985). Mode of action of glyphosate – a literature analysis. In: *The Herbicide Glyphosate*, Grossbard, E. and Atkinson, D (eds), Butterworths, Boston, pp48-74.
- Conner, A.J. (1995). Case study: food safety evaluation of transgenic potato. In: *Application of the Principles of Substantial Equivalence to the Safety Evaluation of Foods or Food Components from Plants Derived by Modern Biotechnology*. Report of a WHO Workshop. World Health Organization, Geneva, pp 23-35.
- Coruzzi, G., Broglie, R., Edwards, C. and Chua, N-H. (1984). Tissue-specific and light regulated expression of a pea nuclear gene encoding the small subunit of ribulose-1,5-bisphosphate carboxylase. *EMBO J.* **3**: 1671-1679.
- Davies, J. *et al* (1986) Aminoglycoside-aminocyclitol antibiotics and their modifying enzymes In: *Antibiotics in laboratory medicine*, 2<sup>nd</sup> ed., Lorian, V., (ed) pp 790-809.
- Davis, B.D., Dulbecco, R., Eisen, H.N. and Ginsberg, H.S. (1980). *Microbiology*, 3<sup>rd</sup> Edition. Harper and Row Publishers, USA, 1355 pp.
- DeBlock, M., Herrera-Estrella, L., Van Montague, M., Schell, J. and Zambryski, P. (1984). Expression of foreign genes in regenerated plants and their progeny. *EMBO J.* **3**: 1681-1689.
- Della-Cioppa, G., Bauer, S.C., Klein, B.K., Shah, D.M., Fraley, R.T. and Kishore G.M., (1986). Translocation of the precursor of 5-enolpyruvylshikimate-3-phosphate synthase into chloroplasts of higher plants *in vitro*. *Proc.Natl.Acad.Sci.* **83**: 6873-6877.
- Depicker, A., Stachel, S., Dhaese, P., Zambryski, P. and Goodman, H.M. (1982). Nopaline synthase: transcript mapping and DNA sequence. *J. Molec. Appl. Genet.* **1**: 561-573.
- Ferré, J. *et al* (1991). *Proc. Natl. Acad. Sci. USA* **88**: 5119-5123.
- Fling, M., Kopf, J. and Richards, C. (1985). Nucleotide sequence of the transposon Tn7 gene encoding an aminoglycoside-modifying enzyme, 3' (9)-O-nucleotidyltransferase. *Nuc. Acids Res.* **13**: 7095-7106.
- Fraley, R.T., Rogers, S.G., Horsch, R.B., Sanders, P.R., Flick, J.S., Adams, S.P., Bittner, M.L., Brand, L.A., Fink, C.L., Fry, J.S., Gallupi, G.R., Goldberg, S.b., Hoffmann, N.L. and Woo, S.C. (1983). Expression of bacterial genes in plant cells. *Proc. Natl. Acad. Sci. (USA)* **80**: 4803-4807.
- Gorbalenya, A.E., Donchenko, A.P., Blinov, V.M. and Koonin, E.V. (1989). Cysteine proteases of positive strand RNA viruses and chymotrypsin-like serine proteases. *FEBS Lett.* **243**: 103-114.
- Habili, N. and Symons, R.H. (1989). Evolutionary relationships between luteoviruses and other RNA plant viruses based on sequence motifs in their putative RNA polymerases and nucleic acid helicases. *Nuc. Acids Res.* **17**: 9543-9555.

Hammond, B.G. and Fuchs R.L. (1998). Safety evaluation for new varieties of food crops developed through biotechnology. In: *Biotechnology and safety assessment*. Thomas JA (ed.), Taylor and Francis, Philadelphia.

Herrnstadt, C., Soares, G.C., Wilcox, E.R. and Edwards, D.L. (1986). A new strain of *Bacillus thuringiensis* with activity against coleopteran insects. *Bio/Technology* **4**: 261-265.

Hofte, H. and Whitely, H.R. (1989). Insecticidal crystal proteins of *Bacillus thuringiensis*. *Microbiol. Rev.* **53**: 242-255.

Horsch, R.B., Fraley, R.T., Rogers, S.G., Sanders, P.R., Lloyd, A. and Hoffmann, N. (1984). Inheritance of functional foreign genes in plants. *Science* **223**: 496-498.

Howe, A.R., Tamayo, F., Brown, S., Armstrong, C., Fromm, M., Hart, J., Padgett, S., Parker, G. and Horsch, R. (1992). Development of glyphosate as a selectable marker for the production of fertile transgenic corn plants. *In Vitro Cell Dev. Biol.* **28**: 124A.

Hubert, P.J. *et al.* (1995). Safety evaluation of transgenic tomatoes expressing *Bt* endotoxin. In: *Application of the Principles of Substantial Equivalence to the Safety Evaluation of Foods or Food Components from Plants Derived by Modern Biotechnology*. Report of WHO Workshop, World Health Organization.

Jones, D.D. and Maryanski, J.H. (1991). Safety considerations in the evaluation of transgenic plants for human food. In: Levin MA and Strauss HS (eds) *Risk assessment in genetic engineering*. New York: McGraw-Hill.

Kaniewski, W., Lawson, C., Loveless, J., Thomas, P., Mowry, T., Reed, G., Mitsky, T., Zalewski, J. and Muskopf, Y. (1995). Expression of potato leafroll virus (PLRV) replicase genes in Russet Burbank potatoes provide field immunity to PLRV. In: *Environmental biotic factors in integrated plant disease control*, Manka, M. (ed), pp 289-292.

Kärenlampi, S. (1996). *Health effects of marker genes in genetically engineered food plants*. Nordic Council of Ministers, Copenhagen, Denmark, 66 pp.

Kemp, A.S., Mellis, C.M., Barnett, D., Sharota, E. and Simpson, J. (1985). Skin test, RAST and clinical reactions to peanut allergens in children. *Clin. Allergy* **15**: 73-78.

Kishore, G. Shah, D., Padgett, S., della-Cioppa, G., Gasser, C., Re, D., Hironaka, C., Taylor, M., Wibbenmayer, J., Eichholtz, D., Hayford, M., Hoffmann, N., Dalannay, X., Horsch, R., Klee, H., Rogers, S., Rochester, S., Brundage, L., Sanders, P. and Fraley, R. (1988). 5-enolpyruvulshikimate-3-phosphate synthase: from biochemistry to genetic engineering of glyphosate tolerance. In: *Biotechnology for Crop Protection*, Knowles, B.H. and Ellar, D.J. (1987). Colloid-osmotic lysis is a general feature of the mechanism of action of *Bacillus thuringiensis*-endotoxins with different insect specificities. *Biochem. Biophys. Acta* **924**: 509-518.

Knowles, B.H. and Ellar, D.J. (1987). Colloid-osmotic lysis is a general feature of the mechanism of action of *Bacillus thuringiensis*-endotoxins with different insect specificities. *Biochem. Biophys. Acta* **924**: 509-518.

Krieg, A., Huger, A.M., Langenbruch, G.A. and Schnetter, W. (1983). *Bacillus thuringiensis* var. *tenebrionis*, a new pathotype effective against larvae of Coleoptera. *Z. Angew. Entomologie* **1983**: 500-508.

Levin, J.G. and Springson, D.B. (1964). The enzymatic formation and isolation of 5-enolpyruvyl shikimate-3-phosphate. *J. Biol. Chem.* **239**: 1142-1150.

Lomonosoff, G.P. (1995). Pathogen-derived resistance to plant viruses. *Annu. Rev. Phytopathol.* **33**: 323-343.

MacIntosh, S.C., Stone, T.B., Sims, S.R., Hunst, P.L., Greenplate, J.T., Marrone, P.G., Perlak, F.J., Fischhoff, D.A. and Fuchs, R.L. (1990). Specificity and efficacy of purified *Bacillus thuringiensis* proteins against agronomically important insects. *J. Invert. Path.* **56**: 258-266.

Mayo, M.A., Robinson, D.J., Jolly, C.A. and Hyman, L. (1989). Nucleotide sequence of potato leafroll luteovirus RNA. *J. Gen. Virol.* **70**: 1037-1051.

McPherson, S., Perlak, F., Fuchs, R., Marrone, P., Lavrik, P. and Fischhoff, D. (1988). Characterisation of the Coleopteran-specific protein gene of *Bacillus thuringiensis* var. *tenebrionis*. *Bio/Technology* **6**: 61-66.

Miller, W.A., Dinesh-Kumar, S.P., and Paul, C.P. (1995). Luteovirus gene expression. *Crit. Rev. Plant Sci.* **14**: 179-211.

Morris, S.C. and Lee, T.H. (1984). The toxicity and teratogenicity of Solanaceae glycoalkaloids, particularly those of potato (*Solanum tuberosum*): a review. *Food Technology in Australia* **36**: 118-123.

Murphy, E.F., True, R.H. and Hogan, J.M. (1967). Detection threshold of sensory panels for mealiness of baked potatoes as related to specific gravity differences. *Amer. Potato. J.* **44**: 442-451.

Murphy, F.A., Fauquet, C.M., Bishop, D.H.L., Ghabrial, S.A., Jarvis, A.W., Martelli, G.P., Mayo, M.A. and Summers, M.D. (eds) (1995). *Virus Taxonomy: Classification and Nomenclature of Viruses*. Sixth Report on the International Committee on Taxonomy of Viruses, Springer-Verlag, New York, pp379-383.

O'Hehir, R.E., Garman, R.D., Greenstein, J.L. and Lamb, J.R. (1991). The specificity and regulation of T-cell responsiveness to allergens. *Ann. Rev. Immunol.* **9**: 67-95.

Pendersen, H.S. and Djortoft, R. (1989). Antigenic and allergenic properties of acidic and basic peptide chains from glycinin. *Food Agric. Immunol.* **1**: 101-109.

Pennington, J.A.T. and Wilkening, V.L. (1997). Final regulations for the nutrition labelling of raw fruits, vegetables, and fish. *J. Amer. Dietetic Assoc.* **97**: 1299-1305.

Perlak, F.J., Deaton, R.W., Armstrong, T.A., Fuchs, R.L., Sims, S.R., Greeplate, J.T. and Fischhoff, D.A. (1990). Insect resistant cotton plants. *Bio/technology* **8**: 939-943.

- Perlak, F.J., Stone, T.B., Muskopf, Y.M., Peterson, L.J., Parker, G.B., McPherson, S.A., Wyman, J., Love, S., Reed, G., Biever, D. and Fischhoff, D.A. (1993). Genetically improved potatoes: protection from damage by Colorado potato beetles. *Plant Mol. Biol.* **22**: 313-321.
- Powell, P.A., Nelson, R.S., De, B., Hoffman, N., Rogers, S.G., Fraley, R.T. and Beachy, R.N. (1986). Delay in disease development in transgenic plants that express the tobacco mosaic virus coat protein gene. *Science* **232**: 738-743.
- Prüfer, D., Tacke, E., Schmitz, J., Kull, B., Kaufmann, A. and Rohde, W. (1992). Ribosomal frameshifting in plants: a novel signal directs the –1 frameshift in the synthesis of the putative viral replicase of potato leafroll luteovirus. *EMBO J.* **11**: 1111-1117.
- Raschke, E., Baumann, G. and Schoffl, F. (1988). Nucleotide sequence analysis of soybean small heatshock protein genes belonging to two different multigene families. *J. Mol. Biol.* **199**: 549-557.
- Richins, R., Scholthof, H. and Shepard, R. (1987). Sequence of figwort mosaic virus DNA (caulimovirus group). *Nucleic Acids Res.* **15**: 8451-8466.
- Sachs, M.I., Jones, R.T. and Unginger, J.W. (1981). Isolation and partial characterisation of major peanut allergen. *J. Allergy Clin. Immunol.* **67**: 27-34.
- Saiki, R.K., Scharf, S., Faloona, F., Mullis, K.B., Horn, G.T., Erlich, H.A. and Arnheim, N. (1985). Enzymatic amplification of  $\beta$ -globin genomic sequences and restriction site analysis for the diagnosis of sickle cell anemia. *Science* **230**: 1350.
- Sanford, J.C. and Johnson, S.A. (1985). The concept of parasite-derived resistance – deriving resistance genes from the parasites own genome. *J. Theor. Biol.* **113**: 395-405.
- Seppälä, U., Alenius, H., Turjanmaa, K., Reunala, T., Palosuo, T. and Kalkkinen, N. (1999). Identification of patatin as a novel allergen for children with positive skin prick test responses to raw potato. *J. Allergy Clin. Immunol.* **103**: 165-171.
- Shaw, K.J., Rather, P.N., Hare, R.S. and Miller, G.H. (1993). Molecular genetics of aminoglycoside resistance genes and familial relationships of the aminoglycoside-modifying enzymes. *Microbiol. Rev.* **57**: 138-163.
- Shibasaki, M., Suzuki, S., Tajima, S., Nemoto, H. and Kuroume, T. (1980). Allergenicity of major component proteins of soybean. *Int. Arch. Allergy Appl. Immunol.* **61**: 441-448.
- Simmonds, N.W. (1976). Potatoes – *Solanum tuberosum* (Solanaceae). In: *Evolution of Crop Plants*, N.W. Simmonds (ed). Longman, London, pp. 279-283.
- Slack, S.A. (1993). Seed certification and seed improvement programs, In: *Potato Health Management*, Rowe, R.C. (ed), APS Press, St Paul, Minnesota.
- Southern, E.M. (1975). Detection of specific sequences among DNA fragments separated by gel electrophoresis. *J. Mol. Biol.* **98**: 503-517.

Stalker, D.M., Thomas, C.M. and Helinski, D.R. (1981). Nucleotide sequence of the region of the origin of replication of the broad host range plasmid RK2. *Mol. Gen. Genet.* **181**: 8-12.

Storey, R.M.J. and Davies, H.V. (1992). Tuber quality. In: *The Potato Crop, The Scientific Basis for Improvement*, P.M. Harris (ed). Chapman and Hall, London.

Sutcliffe, J.G. (1978). Complete nucleotide sequence of the *Escherichia coli* plasmid pBR322. *Symposia on Quantitative Biology* **43**: 77-103.

Taiz, L. and Zeiger, E. (1991). Plant Physiology. The Benjamin/Cummings Publishing Company Inc, California.

Taylor, S.L. Lemanske, R.F. Jr., Bush, R.K. and Busse, W.W. (1987). Food allergens; structure and immunologic properties. *Ann. Allergy* **59**: 93-99.

Taylor, S.L. (1992). Chemistry and detection of food allergens. *Food Technol.* **39**: 146-152.

Taylor, S.L., Nordlee, J.A. and Bush, R.K. (1992). Food allergies. In: *Food Safety Assessment*, Finley, J.W., Robinson, S.F. and Armstrong, D.J. (eds). ACC Symposium Series 484, American Chemical Society, Washington, D.C.

Thomas, P.E. (1997). Sources and dissemination of potato viruses in the Columbia Basin of the Northwestern United States. *Plant Disease* **67**: 744-747.

Van der Wilk, F., Huisman, M.J., Cornelissen, B.J.C., Huttinga, H. and Goldbach, R. (1989). Nucleotide sequence and organisation of potato leafroll virus genomic RNA. *FEBS Lett.* **245**: 51-56.

Van der Wilk, M., Berveek, A.M., Dulleman, J.F. and van den Heuvel, J.M. (1997). The genome-linked protein of potato leafroll virus is located downstream of the putative protease domain of the ORF 1 product. *Virology* **234**: 300-303.

van Gelder, W.W.J. (1990). Steroidal glycoalkaloids: consequences for potato breeding and food safety of utilising wild Solanum species in breeding programmes. In: *Handbook of Natural Toxins, Vol. 6*. (R.F. Keller and A.T. Tu, Eds), Marcel Dekker Inc, New York.

Weidemann, H.L. and Casper, R. (1992). Immunohistological studies on the occurrence of potato leafroll virus in tuber and shoot of the potato plant. *Potato Research* **25**: 99-106.

WHO (1991). Strategies for assessing the safety of foods produced by biotechnology. Report of a joint FAO/WHO Consultation. World Health Organization, Geneva, 59 pp.

WHO (1993). Health aspects of marker genes in genetically modified plants. Report of a WHO Workshop. World Health Organization, Geneva, 32 pp.

Wolfersberger, M.G. (1990). Specificity and mode of action of *Bacillus thuringiensis* insecticidal crystal proteins toxic to lepidopteran larvae: Recent insights from studies utilising midgut brush border membrane vesicles. *Proc. Vth Int. Colloq. Invertebr. Pathol.* August 20-24, 1990, Adelaide, pp. 278-282.

Wong, E.Y., Hironaka, C.M. and Fischhoff, D. (1992). *Arabidopsis thaliana* small subunit leader and transit peptide enhances the expression of *Bacillus thuringiensis* proteins in transgenic plants. *Plant Mol. Biol.* **20**: 81-93.

Zambryski, P. (1992). Chronicles from the *Agrobacterium*-plant cell DNA transfer story. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **43**: 465-490.

## DRAFT REGULATORY IMPACT ASSESSMENT

*Regulatory Impact Assessment*

The Authority is required, in the course of developing regulations suitable for adoption in Australia and New Zealand, to consider the impact of various options (including non-regulatory options) on all sectors of the community, including consumers, the food industry and governments in both countries. The regulatory impact assessment will identify and evaluate, though not be limited to, the costs and benefits of the regulation, and its health, economic and social impacts.

*Identification of affected parties*

1. Governments in Australia and New Zealand
2. Consumers in Australia and New Zealand
3. Manufacturers, producers and importers of food products

*Options**Option 1—To prohibit the sale of food produced using gene technology*

<b>GOVERNMENT</b> Commonwealth, New Zealand Health Departments, State/Territory Health Departments	<b>Benefits</b> <ul style="list-style-type: none"> <li>• no benefits were identified.</li> </ul>	<b>Costs</b> <ul style="list-style-type: none"> <li>• the governments of Australia and New Zealand may be challenged under the WTO to justify the need for more stringent restrictions than apply internationally.</li> <li>• a prohibition on food produced using gene technology in Australia and New Zealand could result in retaliatory trade measures from other countries.</li> <li>• there may be technical problems for AQIS in enforcing such a prohibition at the import barrier.</li> </ul>
<b>INDUSTRY</b> Manufacturers, producers and importers of food products	<b>Benefits</b> <ul style="list-style-type: none"> <li>• Some companies may benefit from being able to exploit niche markets for non-GM products overseas.</li> </ul>	<b>Costs</b> <ul style="list-style-type: none"> <li>• food manufacturers and producers will be unable to use the processed food fractions from foods produced using gene technology thus requiring the switch to non-GM ingredients and the reformulation of many processed food products. The cost to manufacturers of going non-GM has been estimated to be \$A 207m in Australia and \$NZ 37m in New Zealand<sup>5</sup>. This is equivalent to 0.51% of turnover in Australia and 0.19% in New Zealand.</li> </ul>

<sup>5</sup> Report on the costs of labelling genetically modified foods (2000)

<b>CONSUMERS</b>	<b>Benefits</b> <ul style="list-style-type: none"> <li>• no benefits were identified, however as some consumers perceive GM food to be unsafe, they may perceive prohibition of GM food to provide a public health and safety benefit.</li> </ul>	<b>Costs</b> <ul style="list-style-type: none"> <li>• could lead to decreased availability of certain food products.</li> <li>• increased costs to consumers because manufacturers and producers may have to source non-GM ingredients.</li> </ul>
------------------	---	--

*Option 2– to permit the sale of food produced using gene technology*

<b>GOVERNMENT</b> Commonwealth, New Zealand Health Departments, State/Territory Health Departments	<b>Benefits</b> <ul style="list-style-type: none"> <li>• increased innovation and competitiveness in the food industry will benefit the economy.</li> </ul>	<b>Costs</b> <ul style="list-style-type: none"> <li>• minor costs associated with amending the <i>Food Standards Code</i>.</li> </ul>
<b>INDUSTRY</b> Manufacturers, producers and importers of food products	<b>Benefits</b> <ul style="list-style-type: none"> <li>• food producers and manufacturers will be able to capitalise on the latest technology.</li> <li>• food importers will continue to be able to import manufactured products from overseas markets including the USA and Canada where there is no restriction on the use of food produced using gene technology.</li> </ul>	<b>Costs</b> <ul style="list-style-type: none"> <li>• there may be some discrimination against Australian and New Zealand food products in overseas markets that have a preference for non-GM foods (e.g., Japan and the European Union).</li> </ul>
<b>CONSUMERS</b>	<b>Benefits</b> <ul style="list-style-type: none"> <li>• consumers may have access to a greater range of food products.</li> </ul>	<b>Costs</b> <ul style="list-style-type: none"> <li>• those consumers who wish to avoid GM food may experience restricted choice in food products.</li> <li>• those consumers who wish to avoid GM food may have to pay more for non-GM food.</li> </ul>

***Conclusion of the regulatory impact assessment***

Consideration of the regulatory impact for foods produced using gene technology concludes that the benefits of permitting foods produced using gene technology primarily accrue to the government and the food industry, with potentially a small benefit to consumers. These benefits are considered to outweigh the costs to government, consumers and industry, provided the safety assessment does not identify any public health and safety concerns.

## WORLD TRADE ORGANIZATION AGREEMENTS

With the completion of the Uruguay Round of trade negotiations, the World Trade Organization (WTO) was created on 1 January 1995 to provide a forum for facilitating international trade.

The WTO does not engage in any standard-setting activities but is concerned with ensuring that standards and procedures for assessment of and conformity with standards do not create unnecessary obstacles to international trade.

Two agreements which comprise part of the WTO treaty are particularly important for trade in food. They are the;

- Agreement on the Application of Sanitary and Phytosanitary Measures (SPS); and
- Agreement on Technical Barriers to Trade (TBT).

These agreements strongly encourage the use, where appropriate, of international standards, guidelines and recommendations, such as those established by Codex (in relation to composition, labelling, food additives, veterinary drug and pesticide residues, contaminants, methods of analysis and sampling) and the code and guidelines on hygienic practice.

Both Australia and New Zealand are members of the World Trade Organization (WTO) and signatories to the agreements on the Application of Sanitary and Phytosanitary Measures (SPS agreement) and on Technical Barriers to Trade (TBT agreement). Within Australia, the Council of Australian Governments (COAG) has put in place a Memorandum of Understanding binding all States and Territories to the agreements.

The WTO agreements are predicated on a set of underlying principles that standards and other regulatory measures should be:

- based on sound scientific principles;
- developed using consistent risk assessment practices;
- transparent;
- no more trade-restrictive than necessary to achieve a legitimate objective;
- recognise the equivalence of similar measures in other countries; and
- not used as arbitrary barriers to trade.

As members of the WTO both Australia and New Zealand have an obligation to notify the WTO of changes to food standards to enable other member countries of the WTO to make comment. Notification is required in the case of any new or changed standards which may have a significant trade effect and which depart from the relevant international standard (or where no international standard exists). Matters raised in this proposal may be notified to the WTO as either SPS notifications or TBT notifications, or both.

## **SPS Notifications**

These are primarily health related, and refer to any sanitary and phytosanitary measure applied:

- to protect animal or plant life from risks arising from the entry, establishment or spread of pests, diseases or disease carrying organisms;
- to protect human or animal life or health from risks arising from additives, contaminants, toxins or disease-carrying organisms in foods, beverages or foodstuffs;
- to protect human life or health from risks arising from diseases carried by animals, plants or products thereof, or from the entry, establishment or spread of pests; and
- to prevent or limit other damage from the entry, establishment or spread of pests.

The Agreement on the Application of Sanitary or Phytosanitary Measures relates to any sanitary or phytosanitary measure applied to protect animal, plant or human life or health which may directly or indirectly affect international trade. Whether the SPS measure is in the form of a law or mandatory regulation, an advisory guideline, a code of practice or a requirement, it is the purpose of the measure that is important - not its regulatory status. Each WTO member country is entitled to apply SPS measures that are more stringent than the international standards in order to protect the health of its population. In the interests of transparency, each instance of such non-alignment which could result in an impediment to trade must be identified and justified and the documentation of that justification must be readily available

Each member country is also required to apply its methods of risk assessment and management consistently so arrangements under the SPS Agreement do not generate what may really be technical barriers to trade

Under the SPS Agreement, an exporting country can have resort to the WTO's dispute settlement procedures with respect to such a non-alignment. These arrangements mean there is potential for a code of practice to introduce an SPS measure that may bring about non-alignment with international requirements. Such non-alignment would need to be justified scientifically on the grounds that it is necessary to protect human, animal or plant life or health.

## **TBT Notifications**

A technical barrier to trade arises when a mandatory requirement in a country's food regulatory system does not align with the international standard and it is more trade restrictive than is necessary to fulfil a legitimate objective. However, it can be acceptable for a country to have a more stringent requirement than that set internationally for reasons including:

- Maintaining national security;
- Preventing deceptive practices; and
- Protecting human health or safety.

Instances of non-alignment with international standards which could result in trade barriers must be identified and, if questioned, justified. Voluntary codes of practice are not expected to generate technical barriers to trade except where compliance with a code of practice or some aspect of a code of practice is expected. Consequently, it is possible for a voluntary code of practice to be viewed by the WTO as mandatory and subject to all the notification and other provisions applying to mandatory regulations.

The Agreement on Technical Barrier to Trade relates to requirements covering product characteristics or their related processes and production methods. TBT covers measures that are not SPS, such as requirements relating to terminology, symbols, packaging, marking, labelling, food composition and processing methods.

**SUMMARY OF FIRST ROUND PUBLIC SUBMISSIONS FOR APPLICATIONS  
A372, A375, A378, A379, A380, A381, A382, A383, A384, A385, A386, A387 & A388**

**1. National Genetic Awareness Alliance (Aus)**

- believes that the patenting of life-forms and living processes represents a violation of human rights, threat to food security, impediment to medical research and a threat to animal welfare
- believes that current GM techniques are inherently hazardous, and have been shown recently to offer no benefits
  - lower yields with high pesticide input
  - intensification of the corporate monopoly on food
  - spread of antibiotic resistance marker genes and promoter sequences
  - possible increase of allergenicity due to spread of transgenic pollen
- urges governments to use precautionary principle and carry out research into sustainable agricultural methods
- calls for suspension of trials and sale of GM products and public inquiry.

**2. Pola Lekstan and Anna Clements (Aus)**

- are concerned that approval without long-term testing may pose a health threat, that more GM food means less choice for those wanting to avoid it, that Bt may affect non-target organisms, and that herbicide resistance may lead to overuse of chemicals.

**3. Arnold Ward (Aus)**

- questions the system of MRL setting in light of the levels of high glyphosate residues in Roundup Ready soybeans and of other chemicals (including the Bt toxin) in GM crops
- is concerned about detrimental effect of Bt on non-target (beneficial) organisms and on humans, and believes that genetic engineering is imprecise with uncertainties in outcomes
- believes that the concept of substantial equivalence is inadequate and should not be used to avoid more rigorous testing, and that commercial factors are overriding need for basic research. Also believes that ANZFA's arguments defend the needs of biotechnology companies and food processing industry, and that since ANZFA does no testing itself, the results can't be trusted.

**4. Australian GeneEthics Network**

- believes that the data provided is insufficient to make an assessment, and clock should be stopped on the applications. Concerns include:
  - direct health effects of pesticide residues
  - possibility of transfer of antibiotic resistance marker genes leading to resistant bacteria
  - the possibility that transfer of other traits e.g. herbicide tolerance to bacteria, could lead to horizontal spread of unfavourable traits
  - insertion of viral DNA could create new and virulent viruses
  - the possibility that approval could lead to the growing of GMOs in Australia – ecological concerns including effects of, and increases in resistance to, Bt-toxins and the encouragement of increased herbicide use resulting from herbicide-tolerant crops
  - the threat to GE-free status export markets

- believes that the term ‘substantial equivalence’ is not useful– compositional data alone does not establish equivalence

**5. Public and Environmental Health Service (Aus)**

- believes that the data provided should cover both the intentional and unintentional effects of the genetic modification. The unintended consequences of random insertion of new genetic material into the host genome could include loss or change of function of gene or controlling element, dysregulation or amended regulation of the gene or controlling element, or production of a novel hybrid protein which could occur in an unregulated manner. They should also cover any compositional changes e.g. nutrients, anti-nutritional factors, natural toxicants, and define when a change would be considered ‘significant’
- potential effect of introduced proteins on metabolic pathways should be addressed e.g. over-expression or inhibition of enzymes
- data should include details of whether introduced proteins are detectable in whole commodities, processed products and highly processed derivatives
- data should include details of toxicity and allergenicity tests to prove that food is safe, as well as address issues of specificity and potency of proteins. It should also address the ability to support typical growth and well-being
- data for herbicide-tolerant plants should be derived from studies performed on plants treated with herbicide. They should address the human toxicity of the herbicide and whether residues of the herbicide degradation process are present, toxic and/or subject to an MRL.

**6. David Grundy (Aus)**

- considers that the expression of Bt toxins and other chemicals in plant tissues removes the choice of washing chemicals off fruit and vegetables. Believes that Roundup Ready crops have glyphosate or glufosinate molecules genetically attached
- believes that GM crops should not be used for feed given to animals bound for human consumption, that products encouraging antibiotic resistance should not be used, and that labelling should be mandatory for all products containing GM ingredients

**7. Leesa Daniels (Aus) Member of the Genetic Engineering Action Group**

- believes that:
  - scientific research although limited, has brought concerns to light
  - substantial equivalence is a subjective principal
  - comprehensive and mandatory labelling must be urgently implemented
  - the cauliflower mosaic virus (CaMV) promoter could enhance the capability to transfer genes horizontally and has the potential for activating dormant or new viruses
  - antibiotic marker genes could lead to increase in antibiotic resistance
  - several of the transformations encourage the use of pesticides, all of which have shown to be harmful.

**8. Australian Food and Grocery Council**

- fully endorses the policy of minimum affective regulation, supports these applications, and considers that food manufacturers should make their own choice with regard to use of GM crops or products derived from them
- believes that since the growth of GM crops has been approved overseas, they would support their growth in Australia if approved through the GTAC/GMAC/OGTR process

- considers it unfortunate that ANZFA has not negotiated “equivalence” agreements for products already approved overseas to enable approval without having to carry out its own safety assessment. In the absence of such an agreement it supports the ANZFA safety assessment process.
- believes that an appropriate information and labelling scheme would enable consumers to make an informed choice

**9. New Zealand Ministry of Health**

- referred preliminary report to New Zealand Health Research Council, who stated concern that all safety aspects should be carefully considered in the ANZFA process.

**10. Nestle Australia Ltd.**

- supports the continued approval of glufosinate ammonium-tolerant canola, and believes that manufacturers would be disadvantaged were approval not to be granted.

**11. Consumers’ Association of South Australia Inc. & National Council of Women of Australia (CASA supports submission of NCWA)**

- believe that current testing procedure is inadequate and that human trials are the only adequate method, as with testing of new drugs. Also that physiological and neurological effects as well as the toxicological and allergenic effects should be looked at, and that an independent body should be responsible for testing
- do not support the use of antibiotic markers, since they believe they may pose a threat to efficacy of antibiotics in humans
- state that new research has shown that GM soybeans may be a less potent source of phytoestrogens than conventional soybeans confirming the inadequacy of the term ‘substantial equivalence’
- raise the point that although these crops have been approved elsewhere, this situation may change with consumer pressure
- do not accept that it is impossible to source food to ascertain whether or not it contains GM ingredients. Believe that if McCain and Sanitarium can do it, then others should also be able to
- state general concern about the risk that MRLs will be raised as a result of herbicide-tolerant crops being developed, and feel that the calculations used are flawed and are not based on safety criteria
- believe that the use of GM crops in animal feed should also be regulated. A378
- state concern over possible increase in glyphosate use (it is apparently confirmed in one reference that herbicide use increases with herbicide resistant crops), referring to studies that link the chemical to Hodgkin’s lymphoma, and the possibility that Europe may ban it due to adverse effects on beneficial insects. They are particularly concerned that glyphosate is not looked at by the same regulatory body as that looking at GM foods

**A379, A388**

- state concern over the persistence and toxicity of bromoxynil, and consider that these have not been adequately assessed by the US FDA. They understand that the breakdown product of bromoxynil (DBHA) may be more potent than bromoxynil itself, and believe that a safety assessment needs to be done on this too. This is apparently the main residue, and they believe that this may appear in cotton oil and linters.

**A372, A375, A380, A381, A386**

- with respect to glufosinate ammonium, state concern about toxicity, neurotoxicity, teratogenicity and residues in food, soil and water. They believe that Monsanto is

likely to apply for an increase in the MRL, and that such increases are likely to constitute a health hazard

**A380, A382, A383, A384, A385, A386**

- raise issues of adverse effects of Bt toxins on non-target insects and think that it needs more study.

**A387**

- believe that raising the amount of a nutrient in a food may have unknown drawbacks e.g. affecting the efficacy of other nutrients

**12. Health Department of Western Australia**

- highlights various health and environmental concerns:
  - the use of antibiotic resistance genes as markers may transfer resistance to animals via gut bacteria
  - the possibility that microbial gene sequences may contain fragments of other virulent genes, and also that ingesting Bt toxins may be harmful to humans
  - the possibility that insects may be more prone to developing resistance to Bt, since Bt toxins have been found to be released into the soil
- believes that both safety data and gene sequences should be available for public scrutiny

**13. Meat New Zealand**

**A379**

- concerned at how labelling regulations will apply to sausage casings that may contain cotton linters even if they are not to be eaten, i.e. are effectively a processing aid. Think that labelling should only be used to advise the sausage manufacturer not consumers.

**14. BRI Australia**

- supports the approval of all 13 applications provided ANZFA is satisfied with their safety

**15. Food Technology Association of Victoria Inc.**

- supports the approval of all 13 applications provided ANZFA is satisfied with their safety

**16. Diane Davie (Aus)**

- believes all 13 applications should be rejected, since they have not undergone human safety testing here or overseas, and have not been assessed on their ethical merits
- believes that risks include:
  - bacterial and viral vectors which could affect human physiology
  - herbicide and insect-resistance genes, which could increase allergies and antibiotic resistance
  - environmental risks
- also believes that ANZFA must heed the concerns of consumers opposed to GM foods

**17. Martin Hurley, David Hook, Ian Smillie, Margaret Dawson, Tee Rodgers-Hayden, David Lovell-Smith (Natural Law Party), Barbara Brown, Ngaire Mason, Robert Anderson (member, Physicians and Scientists for Responsible Genetics), Louise Carroll, Gilbert Urquart, Caroline Allinson-Dunn, Megan Lewis, Peter Barnes, James Harlow, Gabrielle Dewan, Scott Young, Virginia Murray, Stephanie Chambers, Kay Dyson, Peter Fenwick, Joanne Xerri, Paul True, Josh Gill, James & Peysa Charlwood, Mitta Hirsch, Alan Florence, Nicole Paul, Lawrence Clarke, David Snowman, Reg Paling, Mark and Johanna Blows, David and Bev Semour, Richard and Sharon Moreham (see**

also below), **Stuart Drury and Helen Murphy (All Aus), Brennan Henderson (NZ) – Generic e-mail objection**

- believe that most Australians and New Zealanders do not want GM foods, there are no benefits, and deferral would not be disadvantageous. Approval should be delayed until they are proven safe.
- feel that there is insufficient time to assess these applications thoroughly, and there are so many products under development that there is a high overall risk of a major disaster
- believe that GM foods encourage pesticide use, and applications have made for commercial purposes only, and also that there could be commercial benefit to Australia and New Zealand in remaining GM-free.

**18. Richard and Sharon Moreham (see also above)**

- in addition to the points above, also think that it is unfortunate that the NZ government agreed to joint approval of food, as the Australian public are less educated about the issues surrounding GM foods
- think that approval would only prove that ANZFA serves the interests of large multinational companies rather than those of the public.

**19. Vicky Solah (Aus)**

- is for GM foods if the safety evaluation is carry out using approved, validated methods by an independent body, if the results are made available to consumers, and if all GM food is labelled
- is concerned that transformation may lead to disruption of another gene, and that more research is needed before it is clear whether the process is safe
- with regard to herbicide tolerant crops, is concerned that consumers may not be aware of the need to wash products that have been sprayed, and that this therefore impacts on food safety. Also concerned about environmental impact of these chemicals, and of the possibility of resistance necessitating higher pesticide use in the future.

**20. Dr Rosemary Keighley (Aus)**

- will not purchase foods unless they are certified GM-free. Believes that Australian producers who do not actually use GM products, but who fail to label them as such, will suffer.

**21. Nicola Roil (Aus)**

- believes that GM foods pose health threats and may contaminate non-modified crops

**22. Ian and Fran Fergusson (Aus) – also wrote in the big lot above**

- believe there has been inadequate testing, and are concerned about possible side-effects

**23. Lyndal Vincent (Aus)**

- urges delay of approval until proven safe by extensive testing. Considers that genetic material is being released without knowing what the effects are, and cannot be recalled.
- believes that there is no benefit to the consumer, and that national economic interests are best served by maintaining a GM-free market.

**24. Fay Andary (Aus)**

- does not want any of the 13 products covered by the applications to be approved for inclusion in the food supply

**25. John and Francesca Irving (Aus)**

- thinks that no GE foods should be approved for inclusion in the food chain

**26. Diana Killen (Aus)**

- believes that there is no proven benefit to consumers and in many instances nutritional value is actually lower in GM crops, and it is therefore irresponsible to push through approval without thorough assessment of their long-term safety for public health.
- suggests that research has highlighted adverse allergic reactions and a lowered immune response in some individuals, and that there are health implications with crops designed to be grown with greater concentrations of pesticides
- thinks that labelling is essential for consumers to discriminate in purchasing, and that Australia has a unique opportunity in supply of organic and GM-free food.

**27. Sheila Annesley (Aus)**

- does not want any of the 13 foods included in the food supply.

**28. David and Edwina Ross (Aus)**

- state concern for the future food supplies and well-being of their grandchildren.

**29. Beth Schurr (Aus)**

- wishes to protest against the threat of GM foods, the possible future detrimental effects and the further endangering of the planet.

**30. Beth Eager (Aus)**

- as a parent is concerned that neither the long-term effects on health nor the environment are being considered.

**31. Bruce Pont and Ljiljana Kuzic-Pont (Aus)**

- believe that safety has not been, and cannot be satisfactorily determined, and that any party associated with GM foods could be legally liable should adverse health effects be seen. Thalidomide, smoking, 'Agent Orange' and asbestos all show that such things can affect subsequent generations
- believe that an increase in use of pesticides will result from pesticide-tolerant crops, and that the emphasis should be on organic and/or safe agriculture
- believe that GM-food is a retrograde step, contrary to nature and has the potential to destroy the human race.

**32. Chitta Mylvaganum (Aus)**

- wishes to know what tests were done to assess negative effects on human and environmental health, how thorough they were, what the outcomes were, are the results publicly available, and what further avenues of inquiry are open to the public
- requests the prevention of the import or release of any products until tests are carried out by unbiased scientists in order to prove the lack of health or environmental effects.

**33. John Stevens (Aus)**

- would be concerned if approval were granted before sufficient research had been completed on potential impacts on human health and gene pools of nearby crops. Once grown, spread via pollen would be impossible to stop, and labelling would not prevent exposure by this route
- considers that utmost caution should be exercised and import approval denied indefinitely

**34. Tim Carr (Convenor of the Emergency Committee against GE Foods)**

- believes that GM-foods are produced using a radical and unpredictable new technology so should be subject to more rigorous testing
- states that it is unknown how the introduced gene will interact with and influence genetic expression in the host genome, and could change the chemical nature of the food

- considers that health risks could result from the increased use of pesticides, and also that ANZFA should consider wider environmental, ethical and socio-economic issues.

**35. Jan Kingsbury (Aus)**

- believes that GM-foods could result in loss of economic advantage for Australia and New Zealand since they are known internationally for pure and safe products
- believes that foods are being complicated and pushed by big internationals, and organic farmers are being contaminated by cross-pollination

**36. Teresa Sackett (Aus)**

- believes that:
  - the KPMG report on labelling was prepared in a ridiculously short time and provided limited analysis
  - the proposal of 'no label' for foods which 'may contain' or in which there is 'no evidence' of GM material is inadequate
  - inadequate testing procedures should not be used to declare a product is GM-free just because material can't be detected. In fact testing methods have been developed that can be used to work out the GM content
  - government and industry seem to be favouring the introduction of GM foods. This will result in:
    - (i) increased use of chemicals
    - (ii) destruction of soil life
  - organic farming pay high costs for producing healthy plants, while conventional farmers have little restriction on pollution of air, soil and water. Salinity problems, the death of the Great Barrier Reef, rivers and streams has resulted from ignorance in farming and broader community. Such problems will increase with GM foods.
  - the implication that the public will not understand the issues is wrong. Everyone needs to be fully informed.
- asks the question of whether workers in the food industry are to be better informed, and also why no 'verification documents' are to be required by retailers? Believes that certification schemes should be on a par with those for Kosher foods and organics

**37. John and Sandy Price (Aus)**

- approval of GM foods and seeds should not be allowed, as it is an affront to the sovereignty of Australia and the dignity of the Australian people. The results of the experiment cannot be reversed.

**38. John Scott (NZ)**

- encloses article from The Irish Times, which describes the restrictions that have been placed by the US EPA on the cultivation of GM corn. These appear to have resulted from fears that Bt crops may be harmful to Monarch butterflies and that resistance may develop to Bt

**39. R A Randell (NZ)**

- believes that all GM products should be placed under a moratorium until the Royal Commission of Inquiry has considered the issue, and until all scientific, philosophical, ethical and moral issues have been looked at.

**40. National Council of Women of New Zealand**

- believes that:
  - approval of all 13 applications should be rejected, and that none should be approved for planting.
  - independently-funded body should be responsible for safety assessments

- if it is possible to segregate high-oleic soybeans, then RoundUp Ready soybeans should be segregated too
  - consumers should be made aware of the extent of GM ingredients in their food
  - GM foods, additives or processing aids already on the market must be labelled comprehensively and without extra cost to the consumer – suggest ‘GM unknown’ rather than ‘may contain’
  - appreciates that rejection may contravene the WHO agreement, but consider that the primary role of ANZFA is the assurance of health and safety
- 41. Safe Food Campaign (NZ)**
- believes that approval should be rejected, and a moratorium be put in place until after the Royal Commission of Inquiry, for various reasons:
    - possible effects on non-target insects
    - spread of GM pollen may cause contamination of non-GM (especially organic) crops, and may result in the spread of herbicide-tolerance genes and an increase in resistance development. Cross-pollination is considered a particular risk for canola (A372 & A388). Bt resistance development is noted as being a particular risk for A382, A383 & A384
    - lack of long-term testing means health risks are not known
    - use of broad-spectrum pesticides affects wild flowers and non-target insects.
- 42. Jocelyn Logan, Caroline Phillips (NZ)**
- oppose all 13 applications for the following reasons:
    - testing has not been long-term or independent, precautionary principle should apply. Approval can happen later if GM is proven safe.
    - no clear public benefit, and lack of opportunity for informed choice (immoral and undemocratic). Labelling regulations also unsatisfactory in this respect.
    - environmental concerns (increase in pesticides, threat to organic farming, Bt resistance)
- 43. Robert Anderson (member of Physicians and Scientists for Responsible Genetics - NZ)**
- considers that the GM issue should be reconsidered in the light of the release of internal FDA documents made available for a recent lawsuit aimed at amending their policy. Attached document (presentation given by Steven Druker, Alliance for Bio-integrity) suggests that:
    - scientist’s warnings have been ignored
    - FDA policy may be illegal, violating the Food, Drugs and Cosmetic Act – Mr Druker believes that the term generally-regarded-as-safe (GRAS) cannot apply to foreign DNA
- 44. Stephen Blackheath (NZ)**
- argues that ANZFA’s approach to safety assessments is scientifically unsound:
    - antibiotic resistance marker genes have been cited as being potentially dangerous by groups other than ANZFA e.g. the Royal Society
    - unanticipated toxins and allergens are a concern, and it is suggested that the ANZFA process does not adequately consider these possibilities
    - doesn’t address the question of whether risks exist that are unique to the GM process
    - it relies on data from the manufacturers themselves, with little sway given to evidence from public submissions. Companies have vested interests the

results and cannot be trusted (also gives evidence of Monsanto's past dishonesty)

- believes that ANZFA is subject to undue influence through the directors, and is biased towards being pro-GM
- suggests that RoundUp Ready soybeans are not substantially equivalent as the stems have been found to be more brittle than traditional lines, and may be lower in phytoestrogen content
- also cites the lawsuit being brought by the Alliance for Bio-integrity, and the internal FDA documents that suggest concern from FDA scientists, as evidence of the FDA ignoring important evidence.

**45. Claire Bleakley (NZ)**

- believes that approval should be rejected for various reasons:
  - they may be against Maori views
  - further long-term trials are needed and should be carried out by ANZFA themselves - certain trials have apparently shown effects on immune system, allergies and rare syndromes
  - health concerns of pesticide overuse
  - the possibility of horizontal gene transfer with respect to antibiotic resistance transfer
  - lack of labelling and the use of the unsatisfactory 'substantial equivalence' concept, which makes hazard difficult to assess
  - there is no substantial gain to consumers

## GENERAL ISSUES RAISED IN PUBLIC COMMENTS

The majority of submissions received in response to the Section 14 Gazette Notice, expressed general views against the use of gene technology and asserted that food produced using this technology is unsafe for human. A number of general issues were raised in these submissions and are addressed below.

### 1. *The safety of genetically modified foods for human consumption*

A majority of submitters raised the issue of public health and safety in relation to food produced using gene technology. In particular, it was stated that there has been inadequate testing of genetically modified foods, that there is limited knowledge concerning the risks associated with the technology and that there may be potential long-term risks associated with the consumption of such foods.

- *Evaluation*

It is a reasonable expectation of the community that foods offered for sale are safe and wholesome. In this context, ‘safe’ means that there is a reasonable certainty of no harm. As with other aspects of human activity, the absolute safety of food consumption cannot be guaranteed. Conventionally produced foods, while having a long history of safe use, are associated with human disease and carry a level of risk which must be balanced against the health benefits of a nutritious and varied diet.

Because the use of gene technology in food production is relatively new, and a long history of safe use of these foods has yet to be established, it is appropriate that a cautious approach is taken to the introduction of these foods onto the market. The purpose of the pre-market assessment of a food produced using gene technology under Standard A18 is to establish that the new food is at least as safe as existing foods. The comprehensive nature of the scientific safety assessment, undertaken on a case-by-case basis, for each new modification is reflective of this cautious approach.

The safety assessment focuses on the new gene product(s), including intentional and unintentional effects of the genetic modification, its properties including potential allergenicity, toxicity, compositional differences in the food and its history of use as a food or food product.

Foods produced using gene technology are assessed in part by a comparison with commonly consumed foods that are already regarded as safe. This concept has been adopted by both the World Health Organisation (WHO)/Food and Agriculture Organisation (FAO) and the Organisation for Economic Cooperation and Development (OECD). The Authority has developed detailed procedures for the safety assessment of foods produced using gene technology that are consistent with international protocols developed by these bodies.

## **2. The need for long-term feeding studies**

A number of submissions were concerned about the lack of long-term toxicity studies on genetically modified foods.

- *Evaluation*

Animal studies are a major element in the safety assessment of many compounds, including pesticides, pharmaceuticals, industrial chemicals and food additives. In most cases, the test substance is well characterised, of known purity and of no nutritional value, and human exposure is generally low. It is therefore relatively straightforward to feed such compounds to animals at a range of doses (some several orders of magnitude above expected human exposure levels) in order to identify any potential adverse effects. Establishing a dose-response relationship is a pivotal step in toxicological testing. By determining the level of exposure at which no adverse effects occur, a safe level of exposure for humans can be established which includes appropriate safety factors.

By contrast, foods are complex mixtures of compounds characterised by wide variations in composition and nutritional value. Due to their bulk, they can usually be fed to animals only at low multiples of the amounts that might be present in the human diet. Therefore, in most cases, it is not possible to conduct dose-response experiments for foods in the same way that these experiments are conducted for chemicals. In addition, a key factor to be considered in conducting animal studies on foods is the need to maintain the nutritional value and balance of the diet. A diet that is poorly balanced will compromise the interpretation of any feeding study, since the effects observed will confound and usually override any small adverse effect which may be related to a component or components of the food. Identifying any potentially adverse effects and relating these to an individual component or characteristic of a food can, therefore, be extremely difficult. Another consideration in determining the need for animal studies is whether it is appropriate from an ethical standpoint to subject experimental animals to such a study if it is unlikely to produce meaningful information.

If there is a need to examine the safety of a newly-expressed protein in a genetically-modified food, it is more appropriate to examine the safety of this protein alone in an animal study rather than when it is part of a whole food. For newly-expressed proteins in genetically-modified foods, the acute toxicity is normally examined in experimental animals. In some case, studies up to 14 days have also been performed. These can provide additional re-assurance that the proteins will have no adverse effects in humans when consumed as part of a food. Such experiments can provide more meaningful information than experiments on the whole food. Additional re-assurance regarding the safety of newly-expressed protein can be obtained by examining the digestibility of the new protein in *in-vitro* assays using conditions which simulate the human gastric system.

## **3. Substantial equivalence**

A number of submitters expressed concern regarding the use of the concept of substantial equivalence as part of the assessment process. Some rejected the premise of substantial equivalence on the grounds that differences at the DNA level make foods substantially different.

- *Evaluation*

Substantial equivalence embodies the concept that, as part of the safety assessment of a genetically modified food, a comparison can be made in relation to the characteristics and properties between the new food and traditionally-produced food. This can include physical characteristics and compositional factors, as well as an examination of the levels of naturally occurring allergens, toxins and anti-nutrients.

This allows the safety assessment to focus on any significant differences between the genetically modified food and its conventionally produced counterpart. Genotypic differences (i.e. differences at the DNA level) are not normally considered in a determination of substantial equivalence, if that difference does not significantly change the characteristics for composition of the new food relative to the conventional food.

The concept of substantial equivalence allows for an evaluation of the important constituents of a new food in a systematic manner while, recognizing that there is general acceptance that normally consumed food produced by conventional methods is regarded by the community as safe. It is important to note that, although a genetically modified food may be found to be different in composition to the traditional food, this in itself does not necessarily mean that the food is unsafe or nutritionally inadequate. Each food needs to be evaluated on an individual basis with regard to the significance of any changes in relation to its composition or to its properties.

The concept of substantial equivalence was first espoused by a 1991 Joint Consultation of the Food and Agricultural Organisation (FAO) and the World Health Organisation (WHO) where it was noted that the '*comparison of a final product with one having an acceptable standard of safety provides an important element of safety assessment.*'

The concept has been internationally recognised and embraced as a valuable tool in the safety assessment of foods produced using gene technology. The OECD also advocates an approach to safety assessment based on substantial equivalence as being '*the most practical to address the safety of foods and food components derived through modern biotechnology.*'

#### **4. *The nutritional value of food produced using gene technology***

A small number of submitters expressed concern that the genetic alteration of food decreases its nutritional value.

- *Evaluation*

The assessment of food produced using gene technology by ANZFA entails an exhaustive evaluation of analytical data on any intentional or unintentional compositional changes to the food. This assessment encompasses the major constituents of the food (fat, protein, carbohydrate, fibre, ash and moisture) as well as the key nutrients (amino acids, vitamins, fatty acids). There is no evidence to suggest that genetic modification *per se* reduces the nutritional value of food.

In the future, genetic modification may be used intentionally to improve the nutritional value of food. In this regard, GM foods may be able to assist in addressing the general nutritional needs of the community and also specific dietary needs of sub-populations.

## **5. *Potential toxins and allergens***

Some submitters expressed concerns about the risks of the introduction of new toxins or allergens.

- *Evaluation*

This issue is considered in detail as part of the safety assessment conducted on each new genetic modification applied to a food or commodity crop. New toxins or allergens may be introduced into food by either gene technology or by traditional breeding techniques, or by altered production processes. It is also possible to use these techniques to develop foods specifically where such compounds are significantly reduced or eliminated. One advantage of gene technology, in comparison with these other methods, is that any transferred genes are well characterised and defined, thus the possibility of developing a food with a new toxic or allergenic compound is likely to be reduced.

## **6. *Antibiotic resistance***

Some submitters raised concerns about increased antibiotic resistance resulting from the use of gene technology. Some felt that it would be reassuring if independent biomedical advice were available to reassure the public that the use of antibiotic resistance markers does not pose a risk to the future use of antibiotics in the management of human disease.

- *Evaluation*

The human health considerations in relation to the potential for the development of antibiotic resistance depend on the nature of the novel genes and must be assessed on a case-by case basis. This issue arises because of the use of antibiotic resistance marker genes in the generation of genetically modified plants. In some circumstances, antibiotic resistance genes are linked to the gene of interest, to enable the initial selection of the engineered cells in the laboratory. Those cells that contain the antibiotic resistance marker gene, and hence the gene of interest, will be able to grow in the presence of the antibiotic. Those cells that failed the transformation process are eliminated during the selection procedure.

Concern has arisen that ingestion of food containing copies of antibiotic resistance genes could facilitate the transfer of the gene to bacteria inhabiting the gut of animals and humans. It is argued that these genes may then be transferred to disease causing bacteria and that this would compromise the therapeutic use of these antibiotics.

In 1993, the World Health Organisation Food Safety Unit considered this issue at a Workshop on the health aspects of marker genes in genetically modified plants. It was concluded at that Workshop that the potential for such gene transfers is effectively zero, given the complexity of the steps required. Since this time, several separate expert panels (Report to the Nordic Council, Copenhagen 1996; Advisory Committee on Novel Foods and Processes, UK 1994, 1996; The Royal Society, UK 1998) and numerous scientific papers published in peer reviewed journals have also considered the available evidence on this issue. It is generally agreed that the presence and subsequent transfer of an intact functional gene from transgenic food to micro-organisms in the human intestine is an extremely unlikely event. Furthermore, if this were to occur, bacteria would not normally retain the resistance genes unless there was an

environment for positive selection. The majority of these genes provide for resistance to antibiotics whose use is confined to the laboratory and are not considered to be of major therapeutic use in humans.

Antibiotic resistant bacteria are naturally occurring, ubiquitous and normally inhabit the gut of animals and humans. There is a general consensus that the transfer of antibiotic resistance genes is much more likely to arise from this source and from associated medical practices, rather than from ingested genetically modified food. Even so, at the recent OECD Conference (GM Food Safety: Facts, Uncertainties, and Assessment) held in Edinburgh on 28 February – 1 March 2000, there was general consensus that the continued use of antibiotic marker genes in GM food crops is unnecessary given the existence of adequate alternatives, and should be phased out.

### ***7. Transfer of novel genes***

Some submitters have expressed concern that the transfer of any novel gene may be a health concern.

- *Evaluation*

It is extremely unlikely that novel genetic material will transfer from GM foods to bacteria in the human digestive tract because of the number of complex and unlikely steps that would need to take place consecutively. It is equally unlikely that novel genetic material will transfer from GM foods to human cells via the digestive tract. In considering the potential impact on human health, it is important to note that humans have always consumed large amounts of DNA as a normal component of food and there is no evidence that this consumption has had any adverse effect on human health. Furthermore, current scientific knowledge has not revealed any DNA sequences from ingested foods that have been incorporated into human DNA. Novel DNA sequences in GM foods comprise only a minute fraction of the total DNA in the food (generally less than 0.01%) and are therefore unlikely to pose any special additional risks compared with the large amount of DNA naturally present in all foods.

### ***8. Viral recombination***

Some submitters expressed concern about the long term effects of transferring viral sequences to plants.

- *Evaluation*

This is an issue that is commonly raised because some of the genes that are transferred to plants use a plant virus promoter. Promoters are controlling DNA sequences which act like a switch and enable the transferred genes to be expressed (i.e. to give rise to a protein product) in a plant cell. The routine use of these viral promoters is often confused with research which has shown that plant virus genes, which have been transferred into plants to render them virus-resistant, may recombine with related plant viruses that subsequently infect the plant, creating new viral variants. This research demonstrates that there may be a greater risk to the environment if viral genes are transferred to plants because it may lead to the generation of new plant virus variants capable of infecting a broader range of plants. This is a matter that

will be addressed by the Genetic Manipulation Advisory Committee (GMAC) on a case-by-case basis when it assesses such plants.

However, the presence of plant viruses, plant virus genes or plant virus segments in food is not considered to pose any greater risk to human health as plant viruses are ubiquitous in nature and are commonly found in food eaten by animals and humans. Plant viruses are also biologically incapable of naturally infecting human or animal cells.

### ***9. Labelling of foods produced using gene technology***

A majority of submissions focussed on this issue. Specifically, the submissions called for the labelling of all foods produced using gene technology, regardless of whether they are substantially equivalent to conventional foods. The submitters based their demands for full labelling on the presumption that all foods produced using gene technology are unsafe and on consumer “right to know” arguments. It was stated that full labelling was the only means of identification of foods produced using gene technology available to consumers.

- *Evaluation*

The existing Standard A18 already makes provision for mandatory labelling of genetically modified foods that are substantially different from their conventional counterparts. However, ANZFA is committed to implementing the in-principle decision of ANZFSC Health Ministers of August 1999 to require labelling of all genetically modified foods, including those that are substantially equivalent in composition to the unmodified form. In conjunction with a task force of officials from State and Territory Health Departments and the New Zealand Ministry of Health, ANZFA developed draft revision to Standard A18 in October 1999 that requires labelling of other categories of genetically modified foods. At the Ministers request this draft was circulated for public review and a cost-benefit analysis of full labelling was commissioned. The task force considered both public comments and the cost-benefit analysis in finalising their recommendations to Ministers, which were delivered in May 2000. Ministers are to meet to resolve the issue in July 2000 following whole-of-government consideration of the issue. It is therefore expected that, following a decision and legal amendments to the standard, labelling requirements will be implemented that will apply to all current and subsequent applications.

### ***10. The need for post marketing surveillance of genetically modified foods***

A number of submitters have commented on the need for post-market surveillance of genetically modified food consumption.

- *Evaluation*

Surveillance of potential adverse or beneficial effects of GM foods is seen by many as a logical follow-up to the initial scientific risk assessment. Nevertheless, it is recognised that there are limitations to the application of epidemiology studies, particularly in relation to food components. A key requirement for post-market surveillance systems is that a clear hypothesis be identified for testing. Establishing a system for the surveillance of potential health effects of exposure to novel foods requires monitoring of the consumption patterns of novel foods in the population, and health effects in both “exposed” and “non-exposed” individuals/populations, so that risk estimates can be derived. For any such monitoring

system to be useful, there needs to be a range of exposures, otherwise, any variation in health outcome would be unexplainable by that exposure. Variations in exposure could be apparent over time (temporal trends), space (geographical trends) or both.

Availability of robust data on consumption of the foods in question is vital in order to establish a surveillance system. The other side of the equation is the need for access to data on population health outcomes. Such a system could also be used to identify potential positive health outcomes, such as improved nutritional status or lower cholesterol levels. The availability of linked basic data (e.g. date of birth, sex, geographical location), and the ability to correlate with demographic data, could potentially offer the means of establishing links with food consumption.

The possibility of setting up a post-market health surveillance system for novel foods, including GM foods, has been examined by the UK's Advisory Committee on Novel Foods and Processes (ACNFP). Recognising the many difficulties involved in developing such a system, an initial feasibility study to look at the available data and its usefulness has been proposed. Work is currently being commissioned; when completed in 18 months, it will be subject to peer review. If such a feasibility study suggests that post-market surveillance is practical, methods and details concerning data collection will be determined in the UK, but common strategies might be able to be harmonised internationally in order to minimise the use of resources while maximising the reliability of the final results. This is an area that ANZFA will be monitoring closely, along with international regulatory bodies such as the OECD Taskforce for the Safety of Novel Foods and Feeds.

## **11. *Public consultation and information about gene technology***

A number of submitters were concerned that the public has not been properly consulted or informed by government or ANZFA on the introduction of foods produced using gene technology. Some submitters urged to undertake wider consultation with all affected parties including growers, the food industry and consumers before these food commodities are introduced, and to ensure that adequate consultation is undertaken as part of its assessment process.

- *Evaluation*

The issue of gene technology and its use in food has been under consideration in Australia since 1992. The Agreement between the Governments of Australia and New Zealand for a joint food standard setting system, however, did not occur until 1995, and the New Zealand community therefore had not been consulted on this matter by the Authority until after that time. Consequently, the proposed standard (the current Standard A18) underwent only one round of public comment in New Zealand at which time significant objections were raised by the New Zealand community to the use of gene technology in food production. Many New Zealand consumers, both in these submissions, and in previous submissions to the Authority, have expressed the view that there has been insufficient consultation and a consistent lack of information about gene technology.

Although Standard A18 came into force in May 1999, the public have a continuous and ongoing opportunity to provide comment in relation to applications under the standard. ANZFA's statutory process for all applications to amend the *Food Standards Code* normally involves two rounds of public comment. Furthermore, all the documentation (except for

commercial in confidence information) relating to these applications is available in the public domain, including the safety assessment reports. There is ample evidence that the provision of such information by ANZFA has already significantly stimulated public debate on this matter.

In addition, other government departments including the Environmental Risk Management Authority (ERMA), are potential sources of information about gene technology available to consumers in New Zealand. ERMA is a statutory authority set up by the New Zealand Government to administer the *Hazardous Substances and New Organisms (HSNO) Act 1996*, and has responsibility for assessing the risks to the environment from genetically modified organisms. This body has been assessing applications for the approval of genetically modified organisms since July 1998 and this has involved a number of public meetings.

In response to the concerns raised in public submissions with regard to gene technology and GM foods, ANZFA is in the process of preparing a public discussion paper on the safety assessment process for GM foods. This will be widely available and may assist in addressing some of the concerns raised by the public. Other government and industry bodies are also addressing the broader concerns in relation to gene technology.

## **12. *Maori beliefs and values***

Some New Zealand submitters stated that Maori people find genetic engineering in conflict with their beliefs and values and that, out of respect to Maori, no genetically modified foods should be allowed into New Zealand until a wider discussion, both within Maori and non-Maori, is held.

- *Evaluation*

This issue was also raised during consideration of the proposal for the establishment of Standard A18. At that time, it was stated that the likely implications for Maori regarding genetically modified organisms surround the issues of the rights of Maori to the genetic material from flora and fauna indigenous to New Zealand and the release into the environment of genetically modified organisms. The *HSNO Act 1996* requires that these matters be considered by ERMA.

## **13. *Environmental concerns and the broader regulatory framework***

A number of submitters have raised concerns that genetically modified crops may pose a risk to the environment.

- *Evaluation*

These issues are considered in the assessment processes of GMAC in Australia and the Environmental Risk Management Authority (ERMA) in New Zealand. The Authority does not have the mandate to assess matters relating to environmental risks resulting from the release of food produced using gene technology into the environment. However, links exist between ANZFA and other regulatory agencies in both Australia and New Zealand, and a large degree of information sharing occurs. ANZFA would not recommend the approval of a food produced using gene technology if the genetically modified organism from which it was

derived did not have the appropriate clearance for general release from either GMAC (or its successor) or ERMA, as appropriate.

The regulatory system in Australia will comprise the existing regulators with a legal remit to cover some aspects of GM products (such as imports, food, agricultural and veterinary chemicals):

- the Australia New Zealand Food Authority (ANZFA)
- the Therapeutic Goods Administration (TGA)
- the National Registration Authority for Agricultural and Veterinary Chemicals (NRA)
- the National Industrial Chemicals Notification and Assessment Scheme (NICNAS)
- the Australian Quarantine and Inspection Service (AQIS).

Similarly, various other departments and agencies play their role in the regulatory process in New Zealand:

- the Ministry of Agriculture and Fisheries (MAF)
- the Ministry of Health (MoH)
- the Ministry of Research, Science and Technology (MoRST)

In Australia a new Office of the Gene Technology Regulator (OGTR) will complement the existing arrangements. OGTR will supersede the existing arrangements under the Genetic Manipulation Advisory Committee (GMAC), which advises on research and environmental release of GMOs. OGTR will regulate all GMOs and any 'gap' products (i.e. products for which no other regulator has responsibility).

All GM food is assessed and regulated by the Australia New Zealand Food Authority (ANZFA) under the direction of Commonwealth, State and Territories Health Ministers and the New Zealand Health Minister, sitting as Australia New Zealand Food Standards Council (ANZFSC).

There will be an interface between ANZFA and OGTR. Consequential amendments proposed to the ANZFA Act arising from the draft Gene Technology Bill 2000 will establish a statutory interface between OGTR and ANZFA. This will involve amendments to the ANZFA Act requiring the Authority to advise OGTR of recommendations to ANZFSC regarding the standard for foods produced using gene technology (currently Standard A 18).

#### ***14. Maximum residue levels of agriculture/veterinary chemicals***

A number of submitters have raised concerns that residues of agricultural and veterinary chemicals in genetically modified (e.g. herbicide tolerant) crops may pose a health risk.

- *Response*

Residues of these chemicals can only legally be present if the chemical has been registered for use in Australia and/or New Zealand, and it has been demonstrated that the residue at specified levels does not lead to adverse health impacts. The concentration of a chemical residue that may be present in a food is regulated through maximum residue limits (MRLs). The MRL is the highest residue concentration that is legally permitted in the food. Food

products have to meet the MRL, whether or not they are derived from genetically modified organisms. The MRL does not indicate the chemical residue level that is always present in a food, but it does indicate the highest residue level that could result from the registered conditions of use.

It is important to note that MRLs are not direct public health and safety limits but rather, are primarily indicators of appropriate chemical usage. MRLs are always set at levels lower than, and normally very much lower than, the health and safety limits. The MRL is determined following a comprehensive evaluation of scientific studies on chemistry, metabolism, analytical methods and residue levels. In Australia, the National Registration Authority (NRA) applies to ANZFA to amend the MRLs in the Food Standards Code and the application is considered by ANZFA through its legislated decision making processes. In New Zealand MRLs are set by the Ministry of Health, generally following a request from, and in collaboration with, the Ministry of Agriculture and Forestry. Only following demonstration that the use of agricultural and veterinary chemicals will not result in unsafe residues will the MRL enter into food law, through its inclusion in either the Food Standards Code in Australia, or the New Zealand (Maximum Residue Limits of Agricultural Compounds) Mandatory Food Standard 1999.