

Imported food risk statement

Ready-to-eat cooked crustaceans in which growth of *Listeria monocytogenes* will not occur

Scope: Ready-to-eat (RTE) cooked crustaceans, such as prawns and crabs, in which growth of *Listeria monocytogenes* will not occur. These products have physico-chemical characteristics (pH and water activity) that limit growth of *L. monocytogenes*, e.g. some brined product, and/or are frozen products that are stored and sold frozen to be thawed by the consumer immediately before consumption. RTE cooked crustaceans that are dried or in ambient stable sealed packages are not covered by this risk statement.

Recommendation and rationale

Does *Listeria monocytogenes* in imported RTE cooked crustaceans in which growth of *L. monocytogenes* will not occur present a potential medium or high risk to public health:

Yes

No

Rationale:

- *L. monocytogenes* is a moderately infectious pathogen that can cause severe disease in susceptible populations and is potentially life threatening.
- There have been no reported listeriosis outbreaks associated with RTE cooked crustaceans in which growth of *L. monocytogenes* will not occur, and there is a lack of data on the prevalence of *L. monocytogenes* in these products.
- The cooking step should eliminate the hazard, and although there is potential for post processing contamination, growth of *L. monocytogenes* to levels associated with illness will not occur due to the physico-chemical characteristics of these products.

General description

Nature of the microorganism:

Listeria monocytogenes is a Gram-positive, non-spore forming rod-shaped bacterium that is found throughout the environment. *L. monocytogenes* has been isolated from domestic and wild animals, birds, soil, vegetation, fodder and water; and from the floors, drains and wet areas of food processing factories (FSANZ 2013).

L. monocytogenes is a hardy organism. The temperature range for growth is between -1.5 and 45°C , with the optimal temperature being $30-37^{\circ}\text{C}$ (FSANZ 2013). Temperatures above 50°C are lethal to *L. monocytogenes*, however it can survive for long periods frozen. *L. monocytogenes* will grow in a broad pH range of 4.0–9.6. It can grow at a water activity (a_w) as low as 0.90 and survive for extended periods of time at an a_w of 0.81. *L. monocytogenes* is moderately salt-tolerant, having been reported to grow in 13–14% sodium chloride (Farber et al. 1992; Lado and Yousef 2007). It grows well under both aerobic and anaerobic conditions (Sutherland et al. 2003).

Adverse health effects:

In susceptible populations, *L. monocytogenes* can cause severe disease that is potentially life threatening. People at risk of invasive listeriosis include pregnant women and their foetuses, newborn babies, the elderly and immunocompromised individuals (such as cancer, transplant and HIV/AIDS patients). Less frequently reported, but also at a greater risk, are patients with diabetes, asthma, cirrhosis and ulcerative colitis (FSANZ 2013).

In pregnant women invasive listeriosis can cause spontaneous abortion, stillbirth or neonatal infection. Influenza-like symptoms, fever, and gastrointestinal symptoms can also occur in the mother. In immunocompromised individuals and the elderly, invasive listeriosis can cause potentially fatal bacterial meningitis with symptoms of fever, malaise, ataxia and altered

mental status. The onset of illness of invasive listeriosis generally ranges from 3 days to 3 months after infection. Invasive listeriosis has a fatality rate of 15 – 30% (FDA 2012; FSANZ 2013).

Published data indicate that contaminated food responsible for foodborne illness usually contain levels of *L. monocytogenes* >100 cfu/g (Ryser and Buchanan 2013).

Exposure to *L. monocytogenes* has minimal impact on the general healthy population. If illness does occur it is often mild and may be mistaken for a viral infection or flu (FSANZ 2012).

Consumption patterns:

In the 2011 – 2012 Nutrition and Physical Activity Survey (part of the 2011 – 2013 Australian Health Survey) <1% children (aged 2 – 16 years), 1% of adults (aged 17 – 69 years) and 1% of people aged 70 and above reported consumption of RTE cooked crustaceans (Australian Bureau of Statistics 2011). Mixed foods that contained RTE cooked crustaceans were excluded from the analysis. Survey data was derived from one day of dietary recall data.

Risk factors and risk mitigation

Cooking of RTE crustacean products is a listericidal process. However, re-contamination can occur after this cooking step as *L. monocytogenes* is an ubiquitous organism and can become established in processing environments. Contamination can occur through poor hygienic practices of food handlers or by exposure of product to contaminated air, water, raw materials or food-contact surfaces (Codex 2007).

L. monocytogenes can grow on some cooked crustacean products at refrigeration temperatures (Kataoka et al. 2017; Paranjpye et al. 2008; Parveen et al. 2017). The potential for growth of *L. monocytogenes* is dependent on the characteristics of the food and the conditions and duration of refrigerated storage (FAO/WHO 2004). Growth can be slowed or prevented by modifying the product characteristics through the addition of inhibitory substances or by freezing (Jinneman et al. 2007; Kataoka et al. 2017).

Food preservatives can reduce the growth and survival of *L. monocytogenes* on food products. The effect of preservatives on the growth of *L. monocytogenes* is influenced by temperature, pH, salt content and water activity. For example, sodium nitrite is more effective at preventing growth of *L. monocytogenes* at lower temperatures and pH, or when the sodium chloride concentration is increased (Lado and Yousef 2007).

Several countries, including Australia, permit the use of a specific bacteriophage as a processing aid to eradicate or decrease *L. monocytogenes* on various RTE food products for human consumption, including RTE cooked crustaceans.

The application of HACCP and good hygienic practices in food manufacturing and throughout the supply chain minimise the potential for re-contamination of cooked RTE crustaceans with *L. monocytogenes*. Control measures that prevent the growth of *L. monocytogenes* to high levels in the food are particularly effective at reducing rates of listeriosis, as nearly all cases of listeriosis result from the consumption of high numbers of the pathogen (FSANZ 2013).

The physio-chemical characteristics and the storage conditions of RTE food in which growth of *L. monocytogenes* will not occur are defined in the Australia New Zealand Food Standards Code (the Code)¹. Public information for vulnerable populations to avoid consumption of RTE food, including cooked prawns, that supports the growth of *L. monocytogenes* is available on various government websites including [FSANZ's website](#).

Surveillance information:

Listeria monocytogenes is a notifiable disease in all Australian states and territories with a reported incidence in 2018 of 0.3 cases per 100,000 population (73 cases) which includes both foodborne and non-foodborne cases. The previous five year mean was 0.32 cases per 100,000 population per year (ranging from 0.3 – 0.4 cases per 100,000 population per year) (Department of Health 2019).

Listeriosis associated with consumption of RTE cooked crustaceans in which growth of *L. monocytogenes* will not occur

A search of the scientific literature via EBSCO, US CDC National Outbreak Reporting System and other publications from 2000 to July 2019 did not identify any listeriosis outbreaks associated with consumption of RTE cooked crustaceans in which growth of *L. monocytogenes* will not occur.

Data on the prevalence of *L. monocytogenes* in RTE cooked crustaceans in which growth of *L. monocytogenes* will not occur

A search of the scientific literature via EBSCO and other publications from 2000 to July 2019 did not identify any surveys of the prevalence of *L. monocytogenes* in RTE cooked crustaceans in which growth of *L. monocytogenes* will not occur.

¹ Refer to *Standards or guidelines* section for definition

Standards or guidelines

In Australia:

- Division 2 of [Standard 4.2.1 of the Australia New Zealand Food Standards Code](#) (the Code) states that a seafood business must systematically examine all of its primary production and processing operations to identify potential seafood safety hazards and implement controls that are commensurate with the food safety risk.
- [Standard 1.2.6](#) includes the labelling provisions for the directions for use and the statement of storage conditions.
- Section 1.6.1–4 of [Standard 1.6.1](#) of the Code states:
 - (1) For the purposes of the table to section S27–4, growth of *L. monocytogenes* will not occur in a *RTE food if –
 - (a) the food has a pH less than 4.4 regardless of water activity; or
 - (b) the food has a water activity less than 0.92 regardless of pH; or
 - (c) the food has a pH less than 5.0 in combination with a water activity of less than 0.94; or
 - (d) the food has a refrigerated shelf life no greater than 5 days; or
 - (e) the food is frozen (including foods consumed frozen and those intended to be thawed immediately before consumption); or
 - (f) it can be validated that the level of *L. monocytogenes* will not increase by greater than 0.5 log cfu/g over the food's stated shelf life.
 - (2) For the purposes of the table to section S27–4, a *ready-to-eat food that does not receive a *listericidal process during manufacture is taken to be a food in which growth of *Listeria monocytogenes* will not occur if the level of *Listeria monocytogenes* will not exceed 100 cfu/g within the food's expected shelf life.
- [Schedule 27](#) contains limits for *L. monocytogenes* based on whether growth can occur or not:
 - For RTE food in which growth of *L. monocytogenes* can occur n=5, m=not detected in 25g
 - For RTE food in which growth of *L. monocytogenes* will not occur n=5, m=10² cfu/g
- Under [Standard 1.3.3](#)—Processing aids—of the *Australia New Zealand Food Standards Code*, *Listeria* phage P100 is permitted for use as a processing aid for listericidal treatment of a number of foods, including RTE cooked crustaceans.

Standard 1.6.1 and Schedule 27 mirror the microbiological criteria set out in the Codex guidelines on the application of general principles of food hygiene to the control of *L. monocytogenes* in foods (CAC/GL 61 – 2007) (Codex 2007).

The Codex General principles of food hygiene (CAC/RCP 1 – 1969) follows the food chain from primary production through to final consumption, highlighting the key hygiene controls at each stage (Codex 2003).

Codex Code of practice for fish and fishery products (CAC/RCP 52-2003) covers additional provisions for the growing, harvesting, handling, production, processing, storage, transportation and retail of fish, shellfish and aquatic invertebrates and products thereof from marine and freshwater sources that are intended for human consumption (Codex 2016).

Management approaches used by overseas countries

- European Commission guidelines on microbiological criteria for foodstuffs No 2073/2005 (European Commission 2005) states:
 - For RTE foods in which growth of *L. monocytogenes* can occur the microbiological criterion for *L. monocytogenes* is n=5, c=0, m=absence in 25g (before the food has left immediate control of the food business operator who produced it)
 - For RTE foods in which growth of *L. monocytogenes* cannot occur the microbiological criterion for *L. monocytogenes* is n=5, c=0, m=100 cfu/g
- In New Zealand, RTE crustaceans (lobsters, crabs, bugs, shrimps and prawns and their products) are classed as a food of high regulatory interest, with clearance limits for imported RTE crustaceans of *L. monocytogenes* not detected in 25g (New Zealand Ministry for Primary Industries 2019) .
- United States: zero tolerance in RTE foods (FDA 2015; FSIS 1989).
- Canada: End-product compliance criteria (Health Canada 2011):
 - For RTE foods in which the growth of *L. monocytogenes* can occur; or which have a shelf-life longer than 5 days; or which have not been validated as either Category 2A or 2B: absence in 5x25g samples.
 - Category 2A: RTE foods in which the growth of *L. monocytogenes* can occur but is limited to levels no greater than 100 cfu/g over the course of the stated shelf-life; or RTE foods which have a refrigerated shelf-life of 5 days or less: If shelf life >5 days, processors should validate and verify their processes to ensure that the levels of *L. monocytogenes* are consistently less than or equal to 100 cfu/g throughout the stated shelf-life. If shelf life is ≤5 days, no validation is required.

- Category 2A: RTE foods in which the growth of *L. monocytogenes* cannot occur over the course of the stated shelf life. Such foods are frozen; or meet specified pH and/or water activity parameters. Processors need to monitor their products to ensure that they continue to meet the criteria (e.g., physicochemical parameters such as pH and a_w) that justify their classification.

This risk statement was compiled in: March 2020

References

Australian Bureau of Statistics (2011) National Nutrition and Physical Activity survey, 2011-2012, Basic CURF, CD-ROM. Findings based on ABS CURF data.

Codex (2003) General principles of food hygiene (CAC/RCP 1 - 1969). Codex Alimentarius Commission, Rome. <http://www.fao.org/fao-who-codexalimentarius/codex-texts/codes-of-practice/en/>. Accessed 3 April 2019

Codex (2007) Guidelines on the application of general principals of food hygiene on the control of *Listeria monocytogenes* in foods (CAC/GL 61 - 2007). Codex Alimentarius Commission, Rome. http://www.fao.org/fao-who-codexalimentarius/sh-proxy/pt/?lnk=1&url=https%253A%252F%252Fworkspace.fao.org%252Fsites%252Fcodex%252Fstandards%252FCXG%2B61-2007%252FCXG_061e.pdf. Accessed 23 May 2019

Codex (2016) Code of Practice for Fish and Fishery Products CAC/RCP 52-2003 (amended 2016). Codex Alimentarius Commission, Rome. <http://www.fao.org/fao-who-codexalimentarius/codex-texts/codes-of-practice/en/>

Department of Health (2019) National Notifiable Disease Surveillance Scheme. http://www9.health.gov.au/cda/source/rpt_2.cfm. Accessed 11 January 2019

European Commission (2005) Commission Regulation (EC) No 2073/2005 of 15 November 2005 on microbiological criteria for foodstuffs. Official Journal of the European Union:1–26

FAO/WHO (2004) Risk assessment of *Listeria monocytogenes* in ready-to-eat foods. Food and Agriculture Organization of the United Nations and World Health Organization. Rome. <https://www.who.int/foodsafety/publications/mra4-risk-listeria/en/>. Accessed 3 April 2019

Farber JM, Coates F, Daley E (1992) Minimum water activity requirements for the growth of *Listeria monocytogenes*. Letters in Applied Microbiology 15:103–105

FDA (2012) Bad bug book: Foodborne pathogenic microorganisms and natural toxins handbook., 2nd ed., Silver Spring

FDA (2015) Current good manufacturing practice, hazard analysis, and risk-based preventive controls for human food: (21 CFR Part 117). <https://www.ecfr.gov/cgi-bin/text-idx?SID=85937c272618c7d5c2be5b14b6664f4d&mc=true&node=pt21.2.117&rgn=div5>. Accessed 2 July 2019

FSANZ (2012) *Listeria* and food. Food Standards Australia New Zealand, Canberra. <http://www.foodstandards.gov.au/consumer/safety/listeria/pages/factsheet/listeriaandfoodjuly25590.aspx>. Accessed 3 April 2019

FSANZ (2013) Agents of foodborne illness. Food Standards Australia New Zealand, Canberra. http://www.foodstandards.gov.au/publications/Documents/FSANZ_Foodbornellness_2013_WEB.pdf. Accessed 3 April 2019

FSIS (1989) Revised policy for controlling *Listeria monocytogenes*. <https://www.loc.gov/item/fr054098>. Accessed 2 July 2019

Health Canada (2011) Policy on *Listeria monocytogenes* in Ready-to-Eat Foods. <https://www.canada.ca/en/health-canada/services/food-nutrition/legislation-guidelines/policies/policy-listeria-monocytogenes-ready-eat-foods-2011.html>. Accessed 3 July 2011

Jinneman KC, Wekell MM, Eklund MW (2007) Incidence and behavior of *Listeria monocytogenes* in fish and seafood. In: Ryser E, Marth E (eds) *Listeria, listeriosis and food safety*, 3rd ed, Ch 15. CRC Press Taylor & Francis Group, Boca Raton, pp 617–653

Kataoka A, Wang H, Elliott PH, Whiting RC, Hayman MM (2017) Growth of *Listeria monocytogenes* in thawed frozen foods. *Journal of Food Protection* 80:447–453

Lado B, Yousef AE (2007) Characteristics of *Listeria monocytogenes* important to food processors. In: Ryser E, Marth E (eds) *Listeria, listeriosis and food safety*, 3rd ed, Ch 6. CRC Press Taylor & Francis Group, Boca Raton, pp 157–213

New Zealand Ministry for Primary Industries (2019) Food Notice: Importing Food. <https://www.mpi.govt.nz/dmsdocument/10685>. Accessed 2 April 2019

Paranjpye RN, Peterson ME, Poysky FT, Eklund MW (2008) Incidence, growth, and inactivation of *Listeria monocytogenes* in cooked and peeled cold-water shrimp. *Journal of Aquatic Food Product Technology* 17:266–284

Parveen S, White C, Tamplin ML (2017) A predictive model for the growth of *Listeria monocytogenes* in commercial blue crab (*Callinectes sapidus*). *Journal of Food Protection* 80:1872–1876

Ryser ET, Buchanan RL (2013) *Listeria monocytogenes*. In: Doyle MP, Buchanan RL (eds) *Food microbiology: Fundamentals and frontiers*, 4th ed, Ch 20. ASM Press, Washington DC, pp 503–545

Sutherland PS, Miles DW, Laboyrie DA (2003) *Listeria monocytogenes*. In: Hocking A (ed) *Foodborne microorganisms of public health significance*, 6th. Australian Institute of Food Science and Technology (NSW Branch), Sydney, pp 381–443