



**Effect of Temperature on Phosphomannose Isomerase as Contained in
Test Substance PMI-0105 as Assessed by Enzymatic Activity**

Data requirement:	Not applicable
Author:	Glenn Mims
Date:	Study completed on September 1, 2009
Syngenta Study No.:	PMI-07-05
Report No.:	SSB-023-09
Performing Laboratories:	Syngenta Biotechnology, Inc. Regulatory Science and Trait Safety PO Box 12257 3054 East Cornwallis Rd.

STATEMENTS OF DATA CONFIDENTIALITY CLAIMS


The following statement applies to submissions to the United States Environmental Protection Agency.

Statement of No Data Confidentiality Claims


No claim of confidentiality is made for any information contained in this study on the basis of its falling within the scope of FIFRA section 10(d)(1)(A), (B), or (C).

Company: Syngenta Seeds, Inc. – Field Crops – NAFTA

Company Representative:



Ann Tuttle
Regulatory Affairs Manager



Date

These data are the property of Syngenta Seeds, Inc. – Field Crops – NAFTA and, as such, are considered to be confidential for all purposes other than compliance with the regulations implementing FIFRA Section 10.

Submission of these data in compliance with FIFRA does not constitute a waiver of any right to confidentiality that may exist under any other provision of common law or statute or in any other country.

The following statement applies to submissions to regulatory agencies and other competent authorities other than the U.S. Environmental Protection Agency and all other viewers.

This Document Contains Confidential Business Information

This document contains confidential business information which is proprietary and the publication or disclosure of which would harm the legitimate business interests of the Syngenta group of companies.

The information contained in this document may not be published or disclosed to any third party without the prior consent in writing of the company supplying the relevant information.

The information contained in this document may not be used by any third party including, but not limited to, any regulatory authority to support approval of this product or any other product without the prior consent in writing of the company supplying the relevant information.

STATEMENT CONCERNING GOOD LABORATORY PRACTICES

With the exceptions noted below, this study was conducted in compliance with the relevant provisions of Good Laboratory Practice (GLP) Standards, 40 CFR Part 160 (U.S. EPA 1989) pursuant to the Federal Insecticide, Fungicide, and Rodenticide Act and subsequent revisions.

- The test substance PMI-0105 was characterized under GLP. While the test substance characterization and the enzyme activity reports were not finalized prior to the initiation of this study, they were finalized prior to the completion of this study (Nelson 2008) and (Nelson 2009) respectively.
- The plate imaging software and analysis software were not validated according to the GLPs.

Study Director

Glenn Mims
Glenn Mims
Regulatory Scientist
Regulatory Science and Trait Safety
Syngenta Biotechnology, Inc.

September 1, 2009

Date

Submitted by

Ann Tuttle
Ann Tuttle
Regulatory Affairs Manager
Syngenta Seeds, Inc. – Field Crops – NAFTA
3054 East Cornwallis Rd.
PO Box 12257
Research Triangle Park, NC 27709-2257, USA

August 31, 2009

Date

Representative of the Sponsor

Gerson Graser
Gerson Graser
Team Leader Biochemical Assessment Team
Regulatory Science and Trait Safety
Syngenta Biotechnology, Inc.

August 31, 2009

Date

QUALITY ASSURANCE STATEMENT

Study Title: Effect of Temperature on Phosphomannose Isomerase as Contained in Test Substance PMI-0105 as Assessed by Enzymatic Activity

Study Director: Glenn Mims

Study Number: PMI-07-05

Report Number: SSB-023-09

Pursuant to Good Laboratory Practice Regulations (40 CFR Part 160), this statement verifies that the aforementioned study was inspected and/or audited and the findings reported to Management and to the Study Director by the Quality Assurance Unit on the dates listed below.

<u>Inspection/Audit Type</u>	<u>Inspection/Audit Dates</u>	<u>Reporting Date</u>
Audit Protocol	October 17, 2007	October 17, 2007
In-Progress Inspection	January 18, 2008	January 18, 2008
Final Report Audit (1 st audit)	July 20-21, 2009	July 21, 2009
Final Report Audit (2 nd audit)	July 30, 2009	July 30, 2009
Final Report Audit (3 rd audit)	August 25, 2009	August 25, 2009
Final Report Audit (4 th audit)	August 26, 2009	August 26, 2009

Prepared by: Connie L. Connor

Connie L. Connor
Senior Quality Assurance Auditor
Quality Assurance Unit
Syngenta Biotechnology, Inc.

Date: August 26, 2009

TABLE OF CONTENTS

Statements of Data Confidentiality Claims.....	2
Statement Concerning Good Laboratory Practices.....	3
Quality Assurance Statement.....	4
Table of Contents.....	5
List of Figures.....	6
List of Tables.....	6
List of Acronyms and Abbreviations.....	7
Summary.....	8
Introduction.....	8
Materials and Methods.....	9
Test Substance PMI-0105.....	9
Temperature Experiments and Enzymatic Activity Assays.....	9
Statistical Methods.....	10
Results and Discussion.....	10
Effect of temperature on PMI Enzymatic Activity.....	10
Conclusions.....	12
Records Retention.....	12
Contributing Scientists.....	12
Critical Dates.....	12
References.....	13

LIST OF FIGURES

Figure 1. Effect of Temperature on Enzymatic Activity of PMI as contained in Test Substance PMI-0105.....	11
---	----

LIST OF TABLES

Table 1. Effect of Temperature on Enzymatic Activity of PMI as contained in Test Substance PMI-0105	11
---	----

LIST OF ACRONYMS AND ABBREVIATIONS

loq	limit of quantitation
NADP	β -Nicotinamide adenine dinucleotide phosphate oxidized
NADPH	β -Nicotinamide adenine dinucleotide phosphate reduced
PMI	phosphomannose isomerase
STDEV	standard deviation
Tris	2-amino-2-hydroxymethyl-1,3-propanediol
Tween 20	polyoxyethylene sorbitan monolaurate
U	units - One unit of PMI activity is defined as the amount of enzyme required to catalyze the conversion of 1 μ mol of mannose 6-phosphate to fructose 6-phosphate per min (equivalent to 1 μ mol NADP reduced to NADPH per min) under the described reaction conditions.

SUMMARY

The effect of elevated temperature on phosphomannose isomerase (PMI) protein was determined by incubating solutions of test substance PMI-0105 containing PMI for 30 minutes at temperatures ranging from 25°C to 95°C and subsequently analyzing them for specific phosphomannose isomerase activity. An additional sample of PMI solution was incubated at 4°C to determine a baseline enzymatic activity.

The incubation of PMI at 25°C and 37°C resulted in no loss of enzymatic activity. Incubation at 65°C resulted in loss of enzymatic activity below the enzymatic assay limit of quantitation and the incubation at 95°C resulted in no detectable enzymatic activity.

The data presented in this study support the conclusion that PMI is labile upon heating at temperatures of 65°C and above, as measured by loss of enzymatic activity.

INTRODUCTION

The purpose of this study was to investigate the effect of elevated temperature on the enzymatic activity of phosphomannose isomerase (PMI). The source of PMI in this study was test substance PMI-0105, which was prepared from a recombinant *Escherichia coli* over-expression system prior to the study (Attenborough 2005).

PMI catalyzes the reversible inter-conversion of mannose 6-phosphate and fructose 6-phosphate and has utility as a selectable marker for transformation of many plant species (Bojsen et al. 1994, Joersbo et al. 1998, Negrotto et al. 2000). Plant cells that have been transformed with the *E.coli manA* gene and express PMI protein are able to utilize mannose as a carbon source (Miles and Guest 1984).

In this study PMI, as contained in test substance PMI-0105, was exposed to a range of temperatures (25°C, 37°C, 65°C and 95°C) for 30 minutes. Additionally, the PMI test substance was incubated at 4°C for 30 minutes as a control to determine the baseline PMI activity. Loss or decrease of specific PMI enzymatic activity was used to determine the effect of elevated temperature on the protein.

MATERIALS AND METHODS

Test Substance PMI-0105

Prior to this study test substance PMI-0105 was prepared as described in Attenborough (2005) at Syngenta Protein Science (Jealott's Hill International Research Centre, Bracknell, UK). The *manA* gene was cloned into the inducible, over-expression vector pET-24a in *E. coli* strain BL21(DE3)RP to produce test substance PMI-0105 and has the same amino acid sequence as the endogenous PMI protein produced in *E. coli*. The PMI protein is used as a selectable marker in many of Syngenta's transgenic plant events including 3272 maize.

The lyophilized PMI sample preparation was designated test substance PMI-0105, shipped on dry ice to the Regulatory Science laboratory (Syngenta Biotechnology, Inc.) and stored at $-20^{\circ}\text{C} \pm 8^{\circ}\text{C}$ until further use. Test substance PMI-0105 was found to contain 89.5% PMI by weight (Nelson 2008).

Temperature Experiments and Enzymatic Activity Assays

Test substance PMI-0105 was dissolved in 50 mM Tris (pH 7.0) buffer to a final concentration of 5 μg PMI/ml. Three 1 ml aliquots of the test substance solution were then incubated at each temperature ($25^{\circ}\text{C} \pm 2^{\circ}\text{C}$, $37^{\circ}\text{C} \pm 2^{\circ}\text{C}$, $65^{\circ}\text{C} \pm 2^{\circ}\text{C}$ and $95^{\circ}\text{C} \pm 2^{\circ}\text{C}$) for 30 min. For a baseline control, three additional aliquots were incubated at $4^{\circ}\text{C} \pm 2^{\circ}\text{C}$ for 30 min. After incubation each 1 ml (5 μg PMI/ml) solution was further diluted in 1 ml of 50 mM Tris (pH 7.0) with 0.1% Tween 20 to a concentration of 0.1 ng PMI/ μl .

The enzymatic activity of PMI was monitored for each treatment in triplicate (i.e. three 25°C samples each were aliquoted three times into the assay for a total of nine reactions). The enzymatic reactions were conducted in Corning 96-well plates containing 10 mM mannose 6-phosphate, 1 mM β -nicotinamide adenine dinucleotide phosphate sodium salt hydrate, 2 U/ml phosphoglucose isomerase and 2 U/ml glucose 6-phosphate dehydrogenase in 50 mM Tris buffer (pH 7.0). The reaction was initiated by the addition of 5 ng PMI (50 μl of 0.1 ng/ μl dilution) per well to the assay mixture (total reaction volume 200 μl per well). PMI activity was measured by the production of NADPH in a coupled reaction containing phosphoglucose isomerase (PGI) and glucose 6-phosphate dehydrogenase (G6PDH) based on the method described by Gracy & Noltmann (1968) and Gill et al. (1986) and validated for use in this study by Nelson (2009).



PMI = Phosphomannose isomerase, PGI = Phosphoglucose isomerase, G6PDH = Glucose 6-phosphate dehydrogenase

The production of NADPH was monitored by measuring absorbance increase at 340 nm using a SpectraMax Plus 384 spectrophotometer (Molecular Devices) and SoftMax Pro software. One unit of PMI activity is defined as the amount of enzyme required to catalyze the conversion of 1 μmol of mannose 6-phosphate to fructose 6-phosphate per min (equivalent to 1 μmol NADP reduced to NADPH per min) under the described reaction conditions.

Statistical Methods

Mean values, standard deviations and relative activities were calculated using Microsoft Excel.

RESULTS AND DISCUSSION

No circumstances occurred during conduct of this study that would have adversely affected the quality or integrity of the data generated.

Effect of temperature on PMI Enzymatic Activity

The results shown in Table 1 and Figure 1 indicate that incubation of test substance PMI-0105 at temperatures up to 37°C for 30 minutes did not decrease the enzymatic activity of PMI. Incubation of test substance PMI-0105 at 65°C for 30 minutes resulted in loss of enzymatic activity below the enzymatic assays limit of quantitation, and the incubation at 95°C resulted in no detectable enzymatic activity.

Temperature (°C)	Mean Specific Activity (U ¹ /mg PMI)	STDEV ² (U ¹ /mg PMI)	Relative Activity (%)
4 (control)	420	23	100
25	439	11	105
37	422	11	100
65	<loq ³	<loq ³	<loq ³
95	not detectable ⁴	not detectable ⁴	not detectable ⁴

Table 1. Effect of Temperature on Enzymatic Activity of PMI as contained in Test Substance PMI-0105

¹ One unit of PMI activity is defined as the amount of enzyme required to catalyze the conversion of 1 µmol of mannose 6-phosphate to fructose 6-phosphate per min (equivalent to 1 µmol NADP reduced to NADPH per min) under the described reaction conditions.

²For each temperature three aliquots were incubated and each aliquot was analyzed in triplicate for enzymatic activity as described in MATERIALS AND METHODS under Temperature Experiments and Enzymatic Activity Assays.

³ Less than the limit of quantitation (<loq) for the enzyme assay (i.e. less than 52.2 milli-OD or 6.25 µg NADPH/ml as described in Nelson 2009).

⁴ Not detectable (i.e. below 15.9 milli OD or 1.90 µg NADPH/ml as described in Nelson 2009).




Figure 1. Effect of Temperature on Enzymatic Activity of PMI as contained in Test Substance PMI-0105.

CONCLUSIONS

The effect of temperature on the enzymatic activity of PMI as contained in test substance PMI-0105 was evaluated at various temperatures ranging from 25°C to 95°C in this study. The results indicate that PMI is labile upon incubation for 30 minutes at temperatures of 65°C and higher as shown by loss of enzymatic activity.

RECORDS RETENTION

Raw data, the original copy of this report, and other relevant records are archived at Syngenta Biotechnology, Inc., 3054 East Cornwallis Rd., Research Triangle Park, NC 27709-2257, USA.

CONTRIBUTING SCIENTISTS

The analytical work reported herein was conducted by Glenn Mims B.S. and Andrea Nelson B.S. at Syngenta Biotechnology, Inc.

Reported by: Glenn Mims September 1, 2009
Glenn Mims
Study Director
Regulatory Science and Trait Safety
Syngenta Biotechnology, Inc.
Date

Approved by: Gerson Graser August 31, 2009
Gerson Graser
Team Leader, Biochemical Assessment Team
Regulatory Science and Trait Safety
Syngenta Biotechnology, Inc.
Date

CRITICAL DATES

Study initiation date:	October 18, 2007
Experimental start date:	January 18, 2008
Experimental termination date:	April 7, 2009

REFERENCES

- Attenborough S. 2005. Purification of native PMI from an *E. coli* Expression System (PMI-0105). Syngenta Protein Science Report No. PS-2005-01.
- Bojsen, K., Donaldson, I., Haldrup, A., Joersbo, M., Kreiberg, J. D., Nielsen, J., Okkels, F. T., and Petersen, S. G. 1994. Mannose or Xylose Based Positive Selection. PCT Application Number WO 94/20627.
- Gill, J. F., Vojo, D. and Chakrabarty, M. 1986. Overproduction and Assay of *Pseudomonas aeruginosa* Phosphomannose Isomerase. *J. Bacteriol.* 167:611-615.
- Gracy, R. W and Noltmann, E. A. 1968. Studies on Phosphomannose Isomerase. *J. Biol. Chem.* 243:3161-3168.
- Joersbo, M., Donaldson, I., Kreiberg, J., Petersen, S.G., Brunstedt, J. and Okkels, F.T. 1998. Analysis of mannose selection used for transformation of sugar beet. *Mol. Breeding* 4: 111-117.
- Miles, J. S. and Guest, J. R. 1984. Nucleotide sequence and transcriptional start point of the phosphomannose isomerase gene (*manA*) of *Escherichia coli*. *Gene* 32:41-48.
- Negrotto, D., Jolley, M., Beer, S., Wenck, A.R., Hansen, G. 2000. The use of phosphomannose isomerase as a selectable marker to recover transgenic maize plants (*Zea mays L.*) via *Agrobacterium* transformation. *Plant Cell Reports* 19:798-803.
- Nelson A. 2008. Characterization of Phosphomannose Isomerase Test Substance PMI-0105 and Certificate of Analysis. Syngenta Regulatory Science Report No. SSB-025-07.
- Nelson A. 2009. Single Laboratory Method Validation of Phosphomannose Isomerase (PMI) Enzymatic Activity Assay. Syngenta Regulatory Science Report No. SSB-003-08.
- U.S. EPA. 1989. Good Laboratory Practice Standards. 40 CFR Part 160.