

PSGR

Physicians and Scientists for Global Responsibility

Charitable Trust

Formerly Physicians and Scientists for Responsible Genetics New Zealand

PO Box 8188
TAURANGA 3145

Telephone +64 7 576 5721

5 December 2012

Food Standards Australia New Zealand

Application A1073

Food derived from Herbicide-tolerant Soybean DAS-44406-6:

a transgenic soybean tolerant to 2,4-dichlorophenoxyacetic acid (2,4-D),
glufosinate ammonium and glyphosate.

The Trustees and Members of PSGR urge Food Standards Australia New Zealand to reject this application on the grounds of the facts presented below.

Transgenic soy represents 77% of global soy production. Soy protein is utilised in a multitude of food products¹:

- Traditional soyfoods e.g. tofu, soymilk, soynuts and edamame (green soybeans);
- Food products from veggie burgers to pastas and cereals;
- Dough-based recipes such as pizza;
- Flour isolates and concentrates;
- Baked goods such as breads, cookies, crackers and cakes, doughnuts and pancakes;
- Breakfast cereals – used extensively in hot cereal mixes and breakfast bars;
- Beverages, toppings and dressings, e.g. coffee whiteners, liquid whipped toppings and pre-whipped toppings, sour cream dressings, instant beverages used as meal replacements;
- Processed and whole meat products such as frankfurters, bologna and sausages;
- Dairy analog products include imitation milk and cheese, non-dairy frozen desserts, coffee whiteners and yogurt;
- Soy/milk blends;
- Confectionary;
- Soups, gravies and sauces;
- Canned foods;
- Canned and bottled drinks.

This list demonstrates how easily most consumers would ingest several helpings of soy on a daily basis. With global production at 77%, more than three quarters of these helpings could be of transgenic soy. Estimates suggest up to 80% of US processed food may contain an ingredient from a transgenic crop, such as soy flour or soy lecithin (Hallman et al., 2003).

Because of the regulatory system, consumption would likely be less in New Zealand. Nevertheless, such products do enter the market places in Australia and New Zealand, either as ingredients for the food processing industry, or in imported foods, or in pharmaceutical or dietary supplement products.

In one calculation, assuming 50% of the diet is from transgenic foods and transgenes represent an estimated 0.0005% of the total DNA in food, the consumption figure is 0.5–5 µg/day. While DNA is claimed to be mostly degraded during the industrial process and in the digestive tract, small fragments have been detected in some body tissues such as leukocytes, liver, spleen and gut bacteria (Schubbert et al., 1997). Fragments of orally administered phage M13 and plant DNA have been shown to be taken up by phagocytes as part of their normal function as immune system cells (Schubbert et al., 1998). Fragments could pass into other organs, including the foetus (Beever et al., 2000; Goldstein et al., 2005; Jonas et al., 2001).

In human food crops developed to resist 2,4-D, glufosinate ammonium and glyphosate, consumers will unknowingly be ingesting the resistant transgene/s from whatever part of the plant they consume and will also be exposed to ingesting residues of herbicide applications.ⁱⁱ Whilst the effects of ingesting herbicide-tolerant Soybean DAS-44406-6 may not be as immediate as the effects from spraying, with multiple daily helpings of ingested transgenic soy, cumulative effects are likely to stack up, particularly bearing in mind that other transgenic crops already form part of the human diet. If vested interests have their way, the public will be ingesting food that is near 100% transgenic in time. It is necessary to curb the risks now. It is also necessary for the public to be made aware of the risks, so that they can take any necessary action to avoid food with GE ingredients.

The EC has determined that 1% is an acceptable limit of cross-contamination in non-transgenic products. Consumer interest groups argue only 0% is acceptable. Companies such as Gerber baby foodsⁱⁱⁱ and Frito-Lay^{iv} avoid use of transgenic foods in any of their products to meet public demand. Current technology is unable to detect minute quantities of transgene contamination. Ensuring 0% contamination using existing methodologies cannot be guaranteed. A 1% threshold may even be below current levels of detectability. Even traces so small they are virtually undetectable could have accumulative effects. This is especially true of everyday highly processed food products such as breakfast cereals where the ingredients used to make these products have been pooled from many different sources.

Transgenic foods - human health risks from chemical interactions

We note with concern the absence of data on the potential interactions of chemicals that the product has been designed to resist, and by implication may potentially be used together or applied within a close timeframe.

There is an absence of data to assess potential health risks through unique combinations of chemicals in food that are accepted as probable or feasible. This is an unmanaged risk, and it is important to forestall that risk in the interests of public health and to meet FSANZ's mandated duty of care.

Transgenic food crops and herbicides

Herbicides primarily affect plant metabolism, effectively killing virtually all green plants within a few days of spraying. Resistant transgenic crops are engineered to withstand this spraying. In the process, crops that are contaminated from excessive spraying are created. More spraying has become necessary as resistant weeds increase in number.

The practice of “desiccation” – spraying close to harvest to facilitate easy lifting of the yield - leaves significant concentrations on the harvested crops. Before harvesting, farmers spray crops with broad-spectrum systemic herbicides to kill them off and give them the appearance of uniform maturity. With protein-rich feed the herbicide is sprayed directly onto the grain several days before it is sold as concentrated feed.

The resistant transgenes express in the xylem of plants: leaves, fruit, flowers, pollen, nectar, and guttation fluid of plants.

Glyphosate: The active ingredient in RoundUp is glyphosate. Glyphosate inhibits the enzyme EPSP synthase that is necessary for plants to grow.^v When applied to crops, glyphosate becomes systemic throughout the plant and cannot be removed by washing.

Products containing glyphosate also contain other toxic compounds; e.g. surfactants known as polyoxyethyleneamines (POEA) which can be more toxic than the glyphosate itself. They are irritants of the respiratory tract, eyes and skin and are often contaminated with dioxane, a suspected carcinogen.

A 2009 study ran tests on human cells using formulations of RoundUp that were diluted up to 100,000 times or more. The cells died within 24 hours.^{vi}

A report in the journal *Chemical Research in Toxicology* says the highest maximum residue level (MRL) for glyphosate in food and feed products in the EU is 20 mg/kg. Transgenic soybeans have been found to contain residue levels as high as 17 mg/kg. Malformations in frog and chicken embryos occurred at 2.03 mg/kg, ten times lower than the MRL. Human intake would involve multiple ingestations daily, which could take consumption levels over the MRL.ⁱⁱⁱ

Glufosinate-ammonium inhibits the enzyme glutamine synthetase, necessary for the production of glutamine and for ammonia detoxification. It inhibits the same enzyme in animals.

MAFF UK states that when used as a desiccant, glufosinate residues are detectable in dried peas, field beans, wheat, barley, oilseed rape, and linseed. Wheat grain containing residues ground into flour retained 10-100% of the residue; bran residue levels 10-600% of those in grain.^{vii}

2,4-dichlorophenoxyacetic acid. The ester forms of 2,4-D penetrate foliage and are converted to acid within the plant. This accumulates in cells through passive diffusion, although active transport within the plant may occur. Accumulation is primarily at the meristem tissue of roots and shoots.

2,4-D is rapidly absorbed via oral routes.^{viii} Studies on human volunteers who ingested pure 2,4-D, and on cases of accidental or voluntary acute poisoning with various 2,4-D herbicides, have shown that 2,4-D is very rapidly absorbed from the gut and carried in the blood to cells and tissues throughout the body.^{ix} Human study participants excreted 5 mg/kg 2,4-D in 22.6 hours, 75% in urine within 96 hours. Concentrations in blood plasma paralleled concentration excreted in urine.^x

2,4-D was detected in urine samples collected from all age groups in a large study of the US public. It claimed it is not clear how these residues may affect human health.

2,4-D is a synthetic auxin (plant hormone) and was a major ingredient in Agent Orange, in a 50:50 mixture with 2,4,5-T (trichlorophenoxyacetic acid) in iso-octyl ester form, manufactured for the US Department of Defense primarily by Monsanto Corporation and Dow Chemical. The result of spraying Agent Orange on civilians and military personnel in Vietnam reveals the overt effects of spraying 2,4-D and 2,4,5-T (contaminated with dioxin).

US Vietnam veterans were also affected,^{xi xii} the effects being extensively documented. Presumptive conditions that resulted from the above spraying, and which are currently acknowledged, include: prostate cancer; respiratory cancers; multiple myeloma; type II diabetes; Hodgkin's disease; non-Hodgkin's lymphoma; soft tissue sarcoma; chloracne; porphyria cutanea tarda; peripheral neuropathy; chronic lymphocytic leukaemia; spina bifida in children of veterans exposed to Agent Orange; B cell leukaemia's, such as hairy cell leukaemia; Parkinson's disease; ischemic heart disease and birth defects (Baker, 2010).

Transgenic crops and human health

The World Health Organization states transgenic plants are organisms in which DNA has been altered in such a way that it does not occur naturally. By that description alone, such a plant cannot be “substantially equivalent” to a conventional plant and cannot by any scientific measure be regarded “as safe as the conventional food.”^{xiii}

Regulators continue to increase acceptable residue levels to meet industry demands. EU authorities have further raised the legal limit for glyphosate contamination in wheat and bread to 100 times the legal limit for vegetables and the limit for feed grains 200-fold.

Proponents of genetic engineering claim citizens of the US have eaten transgenic foods for years with no ill effects. This is a seriously misleading statement. Certainly, US citizens have been eating transgenic foods for years, but these foods are unlabelled, there is no mandated registering of potential adverse effects, and there are no substantive independent epidemiological studies on human subjects to see if there are any negative affects to health and wellbeing. Regulators can take note that it took decades to appreciate that trans-fats have caused millions of premature deaths. Lessons can be learned from that experience by applying the precautionary principle to transgenic food crops.^{xiv}

The acknowledged human health risks associated with ingesting transgenes^{xv} are:

Allergenicity: When introducing a novel gene into a plant there is the potential to create a new allergen or cause an allergic reaction in susceptible individuals. For example, engineering Brazil nuts into soybeans was abandoned because of the risk of causing unexpected allergic reactions.^{xvi} Transgenic Starlink™ Corn, approved for animal feed, contaminated the human food chain. The resulting allergic reactions remain controversial. However, a US EPA advisory panel found it was possible Cry9C was an allergen (CDC 2001, Lemaux 2008, Hefle and Taylor 2001), the FDA was unable to rule out allergenicity, experts say it has a “medium likelihood” of being an allergen, and corn in the US was tested until no measurable amounts of StarLink transgenes are determined (EPA 2007). Potentially, minute traces remain.^{xvii}

We know an allergic reaction occurs when ingestion exposes a consumer to a new protein; in the case of transgenic food crops, to a novel protein that may not occur in nature.^{xviii} Substantive studies have failed to find a reason for the recent substantial increases in allergic reactions. These studies include: the International Study of Asthma and Allergies in Childhood (ISAAC)^{xix}, led by Professor Innes Asher of the University of Auckland, a collaborative effort involving over 100 countries; the GA²LEN study^{xx}; and an objective study which showed through serological measurements at three time periods that earlier birth cohorts are less likely to have become atopic than more recent ones^{xxi}.

Medical professionals say the causes of the increased number of allergies in developed countries interplay between genetics and environmental changes. The timeframe is too short to explain a genetic change in the population. Allergies, including food allergies, have increased in New Zealand and other developed countries by epidemic proportions.

New Zealand: A conservative estimate of allergy sufferers in New Zealand is 9000 (2.156%). This low statistic may be because New Zealanders consume fewer transgenes than other countries and because allergies are acknowledged to be infrequently reported.^{xxii}

Australia: One in 20 Australian children suffer from a potentially fatal food allergy. The biggest recent rise is in children under five.^{xxiii}

United States: Allergic disease is the fifth leading chronic disease in the US among all ages, and the third common chronic disease among children under 18 years old. It is estimated that 20% of American children today have allergies.^{xxiv}

United Kingdom: The UK population has the highest prevalence of allergies in Europe and ranks among the highest in the world.^{xxv}

Worldwide: In 2004, the World Allergy Organization's Specialty and Training Council conducted a survey of member societies. Results indicated that in a population of 1.39 billion people 22% may suffer from some form of allergy.^{xxvi}

The above countries have all added transgenes to the daily diet.

Regulators need to consider that a substantial number of the consuming public could express an allergy following ingestion of transgenes, which present proteins unknown in nature.

Unknown effects on human health:

Extracted plant DNA in soil can be taken up in bacteria^{xxvii} and studies have shown transfer between transgenic plant DNA and bacteria can occur. The known mechanisms of transfer are transduction where DNA transfer is mediated by bacteriophages, conjugation where DNA transfer occurs between bacterial cells through conjugation apparatus, and transformation, the uptake of naked DNA. These three processes occur with gastro-intestinal tract bacteria. The most probable method for transfer in the human gut is natural transformation.

No independent substantive studies have been made of the results of feeding transgenes into the human system, excepting one study of the effects of one meal of transgenic soy.^{xxviii} However, the genes and promoters inserted into transgenic plants have characteristics and sequences similar to bacterial genomes and this may increase the likelihood of bacterial expression.

Whereas most DNA is degraded by digestive enzymes in the gut, studies have shown a small percentage survive passage through the gut and would be available to uptake by gut bacteria.^{xxix}

Horizontal gene transfer (HGT) is the transfer of DNA between sexually incompatible organisms and incidences of HGT between bacteria and fungi, between bacteria and protozoa (single-cell organisms), between bacteria and higher plants and animals, and between fungi, and between insects, have been identified.^{xxx} More than 99 percent of soil bacteria cannot be isolated using available culture techniques, which seriously limits detection of HGT. However, most DNA constructs inserted into transgenic crop plants include sections homologous to bacterial DNA. It is accepted DNA homology is an important factor in promoting HGT into bacteria.^{xxxi} DNA transfer can involve DNA carried by a variety of vectors, such as viruses and bacteria, as are used with genetic engineering technology experiments.^{xxxii} The effects of such transfers have not been adequately studied.

Human health and 2,4-D, glufosinate and glyphosate resistant soy

Natural habitats have potential for DNA transfer, e.g. in animal tissues and animal intestines, including those of humans.^{xxxiii xxxiv} Bacterial uptake and expression of DNA can proceed within one minute of ingestion.^{xxxv}

Studies commissioned by the UK Food Standards Agency found transgenic DNA transferred from transgenic soy into microbes in the human intestine after a single meal.^{xxxviii}

Glufosinate-resistant canola/rapeseed was grown in a field trial. Professor Dr Han-Hinrich Kaatz, then Head of Apidology at the Institut für Bienenkunde (Institute for Bee Research) at the University of Jena, now at Martin-Luther-University Halle, Germany, built a netted enclosure in the field that allowed bees to fly freely within it. He installed pollen traps at the beehives to extract pollen samples from the bees' hind legs as they entered the hive. The collected pollen was fed to young honeybees in the laboratory, pollen being their natural diet. After feeding, Professor Kaatz extracted the intestines of young bees and spread the contents on growth medium. He found the gene that confers resistance to glufosinate, the pat-gene, was in the microorganisms, and in some bacteria and in a yeast species. After ingestion, the transgene had been transferred in the bees' gut to the microbes.

Scientists know that bacteria exchange genes and that acquired genes can create pathogenic bacteria. The sequencing of the genome of *E. coli* 0157 showed that 1387 genes had been acquired by HGT. This also showed strains of microbes exist which possess elevated potential to incorporate foreign DNA. For *E. coli* 0157, this potential led to its extreme toxicity.^{xxxvi}

Transgenic technology is designed to replace natural reproductive processes. Selection occurs at the single cell level and the procedure is highly mutagenic, routinely breaching genera barriers. Pleiotropic (unforeseen and unpredictable) effects do occur.^{xxxvii}

A German study found people with no direct contact with agriculture have significant concentrations of glyphosate in their urine. The journal, *Ithaka*, reported that every urine sample collected from city dwellers around Berlin tested positive for glyphosate. Values ranged from 0.5 to 2 nanograms per millilitre (ng/ml); i.e. five to 20 times the permissible upper limit for glyphosate in German drinking water set at 0.1 ng/ml. A conclusion was that glyphosate entered human populations through its presence in daily foods, including glyphosate-resistant soy. The glyphosate would be on resistant transgenic crops and expressed all parts of resistant transgenic plants.^{xxxviii}

Pleiotropic effects can potentially have an unforeseen, negative impact on human health. We know transgenes ingested by human participants found their way into bacteria in the human gut.^{xxvi} Studies on rats show there are appreciable differences in their intestines when fed transgenic potatoes, and other physical aberrations.^{xxxix}

The effects of ingestion of herbicides containing 2,4-D and other compounds, some 2,4-D metabolites or manufacturing by-products were detected in tissues (Geldmacher-Von Mallinckrodt and Lautenbach, 1966; Prescott et al., 1979). Acute high doses administered to laboratory animals produced ataxia, myotonia, and evidence of histological injury to the kidneys, liver, thyroid, eyes, adrenals and gonads.^{xi} The effects of ingesting these contaminants via transgenes have not been adequately studied. Epidemiological studies have reported associations of several types of cancer, such as soft tissue sarcoma and non-Hodgkin's lymphoma, with the exposure to chlorophenoxy herbicides as defoliants or contaminated herbicides.^{xli}

Other risks: It is mandatory for drugs to be identified and monitored for adverse health effects. Without official tracking made of any adverse effects from transgenic foods, it is not easy to identify them when foods or food additives are so widely used. The almost complete lack of labelling of transgenic foods and food ingredients means it is virtually impossible to trace possible allergies or other reactions; and thus easy to dismiss such claims. However, these examples can be drawn on:

- In 2011, doctors at Sherbrooke University Hospital in Quebec, Canada, found Bt-toxin from transgenic corn accumulates in the human body. Researchers found significant levels of the insecticidal protein CryIAb in the blood of pregnant women; CryIAb being present in transgenic Bt crops. The toxin was identified in 93 percent of the pregnant women tested; 80 percent of umbilical blood in their babies; and 67 percent of non-pregnant women.^{xlii}
- Cry9C, also an engineered *Bacillus thuringiensis* (Bt) protein, was engineered into StarLink™ Corn, approved for animal feed but contaminated the human food chain causing adverse reactions. (See also page 4).
- A significant study using human participants showed transgenes can move from transgenic soy into bacteria in the human gut.^{xliii}
- After transgenic soy was introduced in Britain, doctors reported allergic reactions to soy increased 50%.^{xliv} The Irish Doctors' Environmental Association told how increased soy allergies in the Irish Republic mirrored the experience in Britain.^{xlv} The group wants the establishment of a register of diseases thought to be linked to transgenic foods.

Dr Suzanne Wuerthele, a toxicologist and risk assessor, has been a senior scientist at the US Environmental Protection Agency (EPA) for 20 years. Speaking in a personal capacity, she has stated, "The need for careful monitoring is urgent, given the introduction of thousands of GM foods on a global scale..."^{xlvi}

A proponent of the potential benefits of transgenic crops, Ben Mifflin, former director of the Institute of Arable Crops at Rothamsted, near London, concurred: "Under current monitoring conditions, any unanticipated health impact of such foods would need to be a 'monumental disaster' to be detectable."^{xlvi}

Dona and Arvanitoyannis (2009) state: "Most studies with GM foods indicate that they may cause hepatic, pancreatic, renal and reproduction effects and may alter haematological (blood), biochemical, and immunologic parameters, the significance of which remains to be solved with chronic toxicity studies."^{xlvi}

Safety assessments of 2,4-D and DAS-68416-4 soybean

On its website, the US Environmental Protection Agency states that in evaluating pesticides for re-registration it obtains and reviews *a complete set of studies from the producers of the pesticide*, describing the human health and environmental effects of each pesticide (our italics).^{xlix}

The FDA's approval decision^l reads: "*Dow has concluded* that its herbicide tolerant soybean variety, DAS-68416-4 soybean, and the food and feed derived from it are as safe as conventional soybean varieties, and with the exception of the herbicide tolerance traits, are not materially different in composition or other relevant parameters from other soybean varieties now grown, marketed and consumed in the United States. *At this time, based on Dow's data and information, the agency considers Dow's consultation on DAS-68416-4 soybean to be complete*" (our italics).

Neither site gives evidence of independent assessments.

Long-term effects of ingesting transgenes

Scientists and medical professionals do not know what effects there are with humans consuming multiple helpings of transgenic foods daily over long periods of time. Animal studies reveal the potential for conditions presenting now and in the short- and long-term future. Because official bodies accept the word of developers and vested interests continue to deny the possibility of adverse effects, does not mean there are no problems.^{li} A consumer may not know he/she has consumed Botulin toxin - just LD-50 of 0.4 billionth of a gram per kilogram of body weight - until paralysis sets in. Arsenic exploits pathways in cells, binds to proteins, and creates molecular havoc. Small amounts taken over a long period of time produce weakness, confusion and paralysis. Poisons are effective in minuscule amounts, not always undetectable.^{lii}

Recently, the American Academy of Environmental Medicine^{liii} stated: "GM foods pose a serious health risk in the areas of toxicology, allergy and immune function, reproductive health, and metabolic, physiologic and genetic health and are without benefit. There is more than a casual association between GM foods and adverse health effects. There is causation as defined by Hill's Criteria^{liv} in the areas of strength of association, consistency, specificity, biological gradient, and biological plausibility. The strength of association and consistency between GM foods and disease is confirmed in several animal studies."

There is support for the specificity of the association of transgenic foods and specific disease processes. Multiple animal studies show significant immune dysregulation, including upregulation of cytokines associated with asthma, allergy, and inflammation.^{lv}

The Academy^{lv} also says: "... Animal studies also show altered structure and function of the liver, including altered lipid and carbohydrate metabolism as well as cellular changes that could lead to accelerated aging and possibly lead to the accumulation of reactive oxygen species (ROS).^{lvi} Changes in the kidney, pancreas and spleen have also been documented.^{lvii}

"A recent 2008 study links GM corn with infertility, showing a significant decrease in offspring over time and significantly lower litter weight in mice fed GM corn.^{lviii} This study also found that over 400 genes were expressed differently in the mice fed with GM corn. These are genes known to control protein synthesis and modification, cell signalling, cholesterol synthesis, and insulin regulation. Studies also show intestinal damage in animals fed GM foods, including proliferative cell growth^{lix} and disruption of the intestinal immune system.^{lx}

Transgenic food crops are utilised in many forms in human food and animal feed production. All foods potentially present residue. The cumulative effects of humans ingesting transgenic food crops, even in minute amounts, on a daily basis for unlimited periods simply have not been studied. We repeat - it took decades to appreciate that trans-fats have caused millions of premature deaths. The regulatory system should learn from that experience and remove transgenic food crops and feed from the market.^{lxi}

Increasingly, the data show it is biologically possible for transgenic foods to cause adverse health effects in humans.

The valid use of scientific evidence is to set precaution, and not to perpetuate permissive standards for vested interests to use life and Earth's life-support system as one vast laboratory.

PSGR maintains that it is imperative to adopt a precautionary principle approach to transgenic foods.

The Trustees and Members of Physicians and Scientists for Global Responsibility

Paul G Butler, BSc, MB, ChB, Dip.Obst. (Auckland), FRNZCGP
General Practitioner, AUCKLAND

Jon Carapiet, BA(Hons), MPhil.
Senior Market Researcher, AUCKLAND

Bernard J Conlon, MB, BCh, BAO, DCH, DRCOG, DGM, MRCP (UK), FRNZCGP
General Practitioner, ROTORUA

Elvira Dommissie BSc (Hons), PhD, Mus.B, LTCL, AIRMTNZ
Scientist, Crop & Food Research Institute (1985-1993), working on GE onion programme
CHRISTCHURCH

Michael E Godfrey, MBBS, FACAM, FACNEM
Director, Bay of Plenty Environmental Health Clinic, TAURANGA

Elizabeth Harris, MBChB, Dip Obs, CNZSM., CPCH, CNZFP; DMM, FRNZCGP
General Practitioner, KUROW

Frank Rowson BVetMed
Veterinarian, MATAMATA

Peter R Wills, BSc, PhD
Associate Professor, University of Auckland, AUCKLAND

Damian Wojcik, BSc, MBChB, Dip.Theology, Dip.Obst., DCH, FRNZCGP, FIBCMT (USA)
Director and founder of the Northland Environmental Health Clinic, WHANGAREI

Jean Anderson
Businesswoman retired, TAURANGA.

- ⁱ <http://www.soyconnection.com/soyfoods/index.php>.
- ⁱⁱ <http://www.ifrc.org/PageFiles/89755/Photos/307000-WDR-2011-FINAL-email-1.pdf>.
www.gmo-compass.org/eng/agri_biotechnology/gmo_planting/257.global_gm_planting_2009.html.
- ⁱⁱⁱ <http://www.purefood.org/ge/nobabyge.cfm>.
- ^{iv} Frito-Lay's Halfway Measures Banning GE Corn Freak Out Their Competitors: New Seed Planted in Genetic Flap (Washington Post <http://www.purefood.org/ge/fritolayhalf.cfm>).
- ^v Chemical Research in Toxicology 2010; 23(10):1586–1595.
- ^{vi} Chemical Research in Toxicology 2009 Jan; 22(1):97-105.
- ^{vii} <http://www.pan-uk.org/pestnews/Actives/glufosin.htm>. Pers. Comm., MAFF, Pesticides Usage Survey Group. MAFF, York. Agrow No. 273 January 31st 1997, p. 21; Watkins, 1995. MAFF, Evaluation No. 33 : HOE 399866 (Glufosinate-ammonium), Ministry of Agriculture Fisheries and Food, London, 1990. US EPA, Office of Pesticides and Toxic Substances, Experimental Use permit (6340-EUP-RN) and Temporary Tolerance Petition (4G3156) for HOE 39866. Memo from D.S. Saunders to R. Mountfort, Registration Division, 18th April 1985. MAFF, Health and Safety Executive, 1991. Advisory Committee on Pesticides Annual Report 1991, HMSO, London Fujii et al 1996; Watanabe, 1997; Watanabe and Iwasi, 1996. MAFF, Health and Safety Executive, 1991. Advisory Committee on Pesticides Annual Report 1991, HMSO, London; Pesticides Trust [now PAN UK], Crops Resistant to Glutamine Synthetase Inhibitors.
- ^{viii} 2,4-Dichlorophenoxyacetic Acid, CAS No. 94-75-7,
http://www.cdc.gov/exposurereport/data_tables/2.4-DichlorophenoxyaceticAcid_ChemicalInformation.html;
Reregistration Eligibility Decision (RED) 2,4-D: EPA 738-R-05-002. US Environmental Protection Agency, Office of Prevention, Pesticides and Toxic Substances, Office of Pesticide Programmes, 2005. Munro I C et al, Comprehensive, Integrated Review and Evaluation of the Scientific Evidence Relating to the Safety of the Herbicide 2,4-D. J.Am.Call.Toxicol. 1992, 11, (5), 559-664.
<http://npic.orst.edu/factsheets/2.4-DTech.pdf>.
- ^{ix} Curry, 1962; Herlich & Machata, 1963; Nielsen et al., 1965; Dudley & Thapar, 1972; Coutselinis et al., 1977.
- ^x Kohli, J D et al, Absorption and Excretion of 2,4-Dichlorophenoxyacetic Acid in Man, Xenobiotical 1974, 4 (2), 97-100.
- ^{xi} 'Agent Orange: Diseases Associated with Agent Orange Exposure'. Department of Veterans Affairs Office of Public Health and Environmental Hazards, March 25, 2010. York, Geoffrey; Mick, Hayley; 'Last Ghost of the Vietnam War', The Globe and Mail, July 12, 2008. 'Agent Orange blights Vietnam'. BBC News. December 3, 1998. Retrieved April 2010.
- ^{xii} "PL 102-4 and The National Academy of Sciences". nationalacademies.org. 1981-11-03. Retrieved 2012-07-07.
- ^{xiii} World Health Organization (2002) 'Foods derived from modern technology: 20 questions on genetically modified foods';
www.who.int/foodsafety/publications/biotech/20questions/en/index.html.
Society of Toxicology. The safety of genetically modified foods produced through biotechnology; Toxicol. Sci. 2003; 71:2-8.
- ^{xiv} <http://beforeitsnews.com/alternative/2012/09/10-reasons-why-we-dont-need-gm-foods-2466016.html>
<http://www.organicconsumers.org/ge/irish.cfm>.
- ^{xv} <http://www.csa.com/discoveryguides/gmfood/overview.php>.
- ^{xvi} Identification of a Brazil-nut allergen in transgenic soybeans (New England J. of Medicine, Vol 334, No 11, pp 688-692, 1996).
- ^{xvii} <http://academicreview.org/reviewed-content/genetic-roulette/section-3/3-5-starlink/> CDC (2001). Investigation of Human Health Effects Associated with Potential Exposure to Genetically Modified corn. A Report to the US Food and Drug Administration from the Centres for Disease Control and Prevention, June 11, 2001, www.cdc.gov/nceh/ehhe/Cry9cReport/ accessed 7 January 2009. EPA (2007). US Environmental Protection Agency Office of Pesticide Programs (Draft White Paper) Concerning Dietary Exposure To Cry9c Protein Produced By Starlink® Corn And The Potential risks Associated With Such Exposure October 16, 2007. Goodman RE., Vieths S, Sampson HA, Hill D, Ebisawa M, Taylor, SL, van Ree, R. 2008. 'Allergenicity assessment of genetically modified crops: what makes sense?' Nat. Biotech. 26: 73-81. Hefle S and Taylor S (2001). 'Will genetically modified foods be allergenic?' J Allergy Clin Immunol. 107: 765-771. Lemaux P (2008). 3.9. Were Foods Made From Bt Corn Removed from the Market Because of Allergenicity Concerns? In Genetically engineered plants and foods: a scientist's analysis of the issues (Part I). Annual Review Plant Biology 59:771–812. Petersen BJ, Rachman NJ, Watters JL (2001). Estimated potential dietary intake of cry9c protein based on measurements of cry9c in processed foods made from 100% Starlink™ corn Aventis Corporation filing with EPA, April 21, 2001. Shillito RD, MacIntosh S, Kowite WJ. "Detection of Cry9C protein in dry milled, wet milled and masa processed fractions and processed foods made from 100% StarLink™ grain." Aventis Corporation filing with EPA, April 17, 2001. Tomaselli KP, Allergic reaction: Food allergies increasing, especially among children. American Medical News (AMA) On-line January 5, 2009. www.ama-assn.org/amednews/2008/12/29/hlsa1229.htm.
- ^{xviii} <http://science.nationalgeographic.com/science/health-and-human-body/human-body/allergy-misery-article.html>.
- ^{xix} Lai C, Beasley R, Crane J. et al. Global variation in the prevalence and severity of asthma symptoms: Phase Three of the ISAAC study. Thorax 2009; 64: 476-483.
- ^{xx} Allergy 2009; 64: 969-977; Pirkko Mattila et al, Clinical and Translational Allergy 2011, 1:5 doi:10.1186/2045-7022-1-5, <http://www.ctajournal.com/content/1/1/5>.
- ^{xxi} Law M et al. Changes in Atopy over a quarter of a century, based on cross-sectional data at three time periods. BMJ Vol 330 21 May 2005.
- ^{xxii} www.beehive.govt.nz/release/food. <https://www.cia.gov/library/publications/the-world-factbook/print/nz.html>.
- ^{xxiii} (J of Allergy and Clinical Immunology (2007;120:878-84). www.healthbrands.com.au/topics/Kids/news/.
www.foodallergy.org/featuredtopic.html. www20.sbs.com.au/podcasting/index.php?action=feeddetails&feedid=53&id=7961 .
- ^{xxiv} www.aaaai.org/media/resources/media_kit/allergy_statistics.stm. <http://foodallergyawareness.com/>.
www.foodallergy.org/media/press_releases/anaphylaxisstudy.html. www.allergykids.com/index.php?id=3. Statistics from <http://allergicare.com/>.
- ^{xxv} www.aaircharity.org/allergies/oralallergy.htm. Statistics from <http://allergicare.com/>.
- ^{xxvi} Statistics from <http://allergicare.com/>.
- ^{xxvii} Gebhard F. and Smalla K. (1999) Monitoring field releases of genetically modified sugar beets for persistence of transgenic plant DNA and horizontal gene transfer, FEMS Microbiology Ecology, 28(3).

- xxviii Netherwood T. et al. (2001), Transgenes in genetically modified Soya survive passage through the human small bowel but are completely degraded in the colon, Food Standards Agency Project GO1008.
- xxix 'Summary of current scientific awareness of the effect of genetically modified organisms on the natural environment', Report prepared for the Ministry of the Environment by Glare, T R, O'Callaghan, M, Malone, L A and Burgess E P J.
www.mfe.govt.nz/publications/organisms/scientific-awareness-jun01.pdf.
- xxx Ho M.W., Department of Biology, UK Open University, Excerpt from a talk given at The National Council of Women of Great Britain Symposium on Food: Facts, Fallacies and Fears, 22 March 1996, Darlington. Gebhard F and Smalla K, (1999), 'Monitoring field releases of genetically modified sugar beets for persistence of transgenic plant DNA and horizontal gene transfer', FEMS Microbiology Ecology 28, 261-272.
- xxxi Flint H. et al. (2001), Survival of ingested DNA in the gut and the potential for genetic transformation of resident bacteria, Food Standards Agency Project FSG01007.
- xxxii Traavik T, Professor and Scientific Director at the Norwegian Institute of Gene Ecology, Department of Virology, University of Tromsø, Norway, Evidence given at the trial of 28 Greenpeace activists against AgrEvo (92 references cited), April 2000.
- xxxiii Doerfler W. and Schubert R. (1998) Uptake of foreign DNA from the environment: the gastrointestinal tract and the placenta as portals of entry. Wien Klin. Wochenschr. 110 (2).
- xxxiv Traavik T, Professor and Scientific Director at the Norwegian Institute of Gene Ecology, Department of Virology, University of Tromsø, Norway, Evidence given at the trial of 28.
- xxxv Flint H. and Scott K. (2001), Dissemination of GM DNA and antibiotic resistance genes via rumen micro-organisms, Food Standards Agency Project G01011.
- xxxvi Perna N.T. et al. (2001) Genome sequence of enterohaemorrhagic *Escherichia coli* O157:H7 Nature 409.
- xxxvii Freese W, Schubert D. Safety testing and regulation of genetically engineered foods. Biotechnology and Genetic Engineering Reviews. Nov 2004. 21.
- xxxviii <http://www.ithaka-journal.net/herbicide-im-urin?lang=en>.
- xxxix Effect of diets containing genetically modified potatoes expressing *Galanthus nivalis* lectin on rat small intestine (Lancet, Vol 354, No 9187, pp 1353-1354, Oct 1999). Safety of genetically modified food questioned: Interview with gene scientist, Dr Arpad Pusztai (www.wsos.org/articles/1999/jun1999/gmo-j03.shtml). Science, Vol 286, p 656, Oct 1999.
- xl (NTP, 2006; U.S. EPA, 2005).
- xli Garabrant and Philbert, 2002; IOM, 2003; IPCS, 1996; Pearce and McLean; 2005; U.S. EPA, 2005).
- xlii Aris A, Leblanc S, Reproductive Toxicology. 2011; 31(4), www.ncbi.nlm.nih.gov/pubmed/21338670.
- xliii Netherwood T et al, Nat Biotechnol. 2004 Feb;22(2):204-9, www.ncbi.nlm.nih.gov/pubmed/14730317.
- xliv www.responsibletechnology.org/gmo-dangers/health-risks/articles-about-risks-by-jeffrey-smith/Genetically-Engineered-Foods-May-Cause-Rising-Food-Allergies-Genetically-Engineered-Soybeans-May-2007.
- xlvi <http://www.netlink.de/gen/Zeitung/1999/990312.htm>.
- xlvi www.nature.com/nbt/journal/v23/n2/full/nbt0205-170.html. Nature Biotechnology 23, 170 (2005) doi:10.1038/nbt0205-170.
- xlvi Nature, Volume 398:651 April 22, 1999 www.natural-law.ca/genetic/NewsJuly-Aug99/GEN8-15NaturUkUprootUSA.html.
- xlvi Dona A, Arvantoyannis I S, Health Risks of genetically modified foods. Crit Rev Food Sci Nutri. 2009; 49: 164-175.
http://www.somloqueseembrem.org/img_editor/file/Dona&Arvanitoyannis2009.pdf.
- xlvi 30 June 2005, EPA-738-F-05-002, Pesticide Reregistration http://www.epa.gov/oppsrrd1/REDs/factsheets/24d_fs.htm.
- l Biotechnology Consultation Note to the File BNF No. 000124, 8 November 2011; Subject: DAS-68416-4, herbicide tolerant soybean, <http://www.fda.gov/Food/Biotechnology/Submissions/ucm283173.htm>.
- li The US Environmental Protection Agency on 2,4-D www.epa.gov/oppsrrd1/REDs/factsheets/24d_fs.htm.
- lii <http://www.straightdope.com/columns/read/2707/whats-the-fastest-acting-most-lethal-poison>.
- lii <http://www.aemonline.org/gmopost.html>.
- lii Hill, AB. The environment and disease: association or causation? Proceeding of the Royal Society of Medicine 1965; 58:295-300.
- lii Finamore A, Roselli M, Britti S, et al. Intestinal and peripheral immune response to MON 810 maize ingestion in weaning and old mice. J Agric. Food Chem. 2008; 56(23):11533-11539. Kroghsbo S, Madsen C, Poulsen M, et al. Immunotoxicological studies of genetically modified rice expression PHA-E lectin or Bt toxin in Wistar rats. Toxicology. 2008; 245:24-34.
- lii Malatesta M, Boraldi F, Annovi G, et al. A long-term study on female mice fed on a genetically modified soybean: effects on liver ageing. Histochem Cell Biol. 2008; 130:967-977. Velimirov A, Binter C, Zentek J. Biological effects of transgenic maize NK603xMON810 fed in long term reproduction studies in mice. Report-Federal Ministry of Health, Family and Youth. 2008.
- lii Kilic A, Aday M. A three generational study with genetically modified Bt corn in rats: biochemical and histopathological investigation. Food Chem. Toxicol. 2008; 46(3):1164-11707
- lii Finamore A, Roselli M, Britti S, et al. Intestinal and peripheral immune response to MON 810 maize ingestion in weaning and old mice. J Agric. Food Chem. 2008; 56(23):11533-11539. Velimirov A, Binter C, Zentek J. Biological effects of transgenic maize NK603xMON810 fed in long term reproduction studies in mice. Report-Federal Ministry of Health, Family and Youth. 2008.
- lii Kilic A, Aday M. A three generational study with genetically modified Bt corn in rats: biochemical and histopathological investigation. Food Chem. Toxicol. 2008; 46(3):1164-1170.
- lii Velimirov A, Binter C, Zentek J. Biological effects of transgenic maize NK603xMON810 fed in long term reproduction studies in mice. Report-Federal Ministry of Health, Family and Youth. 2008.
- lii Ewen S, Pusztai A. Effects of diets containing genetically modified potatoes expressing *Galanthus nivalis* lectin on rat small intestine. Lancet. 354:1353-1354.
- lii Finamore A, Roselli M, Britti S, et al. Intestinal and peripheral immune response to MON 810 maize ingestion in weaning and old mice. J Agric. Food Chem. 2008; 56(23):11533-11539.
- lii <http://beforeitsnews.com/alternative/2012/09/10-reasons-why-we-dont-need-gm-foods-2466016.html>
<http://www.organicconsumers.org/ge/irish.cfm>.